



The Australian Prevention
Partnership Centre
Systems and solutions for better health

Secondary prevention of chronic pain

Rapid review and mapping of options for
Primary Health Networks

October 2019

Secondary prevention of chronic pain

Rapid review and mapping of options for Primary Health Networks

Prepared by: Menzies Centre for Health Policy, Faculty of Medicine and Health, University of Sydney for The Australian Prevention Partnership Centre

Contributing authors: Dr Simone De Morgan, Professor Fiona Blyth, Ms Leah Marks, Dr Duncan Sanders, Dr Manasi Mittinty, Professor Michael Nicholas

© Sax Institute 2019



All material and work produced by the Sax Institute is protected by copyright. The Institute reserves the right to set terms and conditions for any use of this material. This product, excluding the Institute's logo and associated logos, and any material owned by third parties, is made available under a Creative Commons Attribution–NonCommercial–ShareAlike 4.0 International licence.

You are free to copy and redistribute the material in any medium or format, provided you attribute the work to the Sax Institute, acknowledge that the Sax Institute owns the copyright, and indicate if any changes have been made to the material. You may not use the material for commercial purposes. If you remix, transform or build upon the material, you must distribute your contributions under the same licence as the original.

Enquiries about any use of this material outside the scope of this licence can be sent to:
preventioncentre@saxinstitute.org.au

Suggested citation: De Morgan S, Blyth F, Marks L, Sanders D, Mittinty M, Nicholas M. Secondary prevention of chronic pain: rapid review and mapping of options for Primary Health Networks. The Australian Prevention Partnership Centre and the University of Sydney, October 2019.



Australian Government
Department of Health

Funding for this research has been provided from the Australian Government's Medical Research Future Fund (MRFF). The MRFF provides funding to support health and medical research and innovation, with the objective of improving the health and wellbeing of Australians. MRFF funding has been provided to The Australian Prevention Partnership Centre under the MRFF Boosting Preventive Health Research Program. Further information on the MRFF is available at www.health.gov.au/mrff

Disclaimer: This evidence review is not a comprehensive review of all literature relating to the topic area. It was current at the time of production (but not necessarily at the time of publication) and is based on sources believed to be reliable.

Contents

Executive summary	5
Chronic Pain Project	5
Phase 1 of the Chronic Pain Project	6
Purpose and scope of the rapid review	6
Principles for the secondary prevention of chronic pain	7
Mapping of the options for Primary Health Networks	9
Resources for consumers and health professionals	13
Conclusion	13
1. Rationale for the secondary prevention of chronic pain	14
2. Principles for the secondary prevention of chronic pain	17
3. Mapping of the options for Primary Health Networks with supporting evidence	24
3.1 Options related to Goal 1 (consumer and community initiatives)	28
3.2 Options related to Goal 2 (health professional capacity building)	51
3.3 Options related to Goal 3 (health systems support initiatives)	62
4. Consumer resources relevant to the secondary prevention of chronic pain	69
5. Health professional education and training and other resources relevant to the secondary prevention of chronic pain	71
References	77
Appendix 1: Chronic Pain Project Steering Committee	83
Appendix 2: Scope of the rapid review	84
Appendix 3: Evidence map of the options related to Goal 1	88
Appendix 4: Evidence map of the options related to Goal 2	96
Appendix 5: Evidence map of the options related to Goal 3	100
Appendix 6: Enablers to implementing consumer pain programs identified in Phase 1 of the Chronic Pain Project	103
Appendix 7: Enablers to implementing education and training initiatives identified in Phase 1 of the Chronic Pain Project	104
Appendix 8: Enablers to implementing HealthPathways identified in in Phase 1 of the Chronic Pain Project	105
Appendix 9: Potential prototype of a consumer pain program for people with (sub)acute pain who are at risk of developing chronic pain	106
Appendix 10: Key content of the training packages in the IMPaCT Back trial	115
Appendix 11: Content of a GP trainee education initiative about opioids implemented in Australia	117

Figures

Figure 1: Principles for the secondary prevention of chronic pain.....	17
Figure 2: Key elements of the principles for the secondary prevention of chronic pain.....	18
Figure 3: Risk factors for the development of chronic postsurgical pain	20
Figure 4: Risk factors for low back pain and associated disability.....	21
Figure 5: Yellow flags.....	22

Tables

Table 1: Map of the options for PHNs related to the secondary prevention of chronic pain.....	25
Table 2: Evidence map of the options related to Goal 1	88
Table 3: Evidence map of the options related to Goal 2	96
Table 4: Evidence map of the options related to Goal 3	100

Executive summary

Chronic Pain Project

Chronic pain is a substantial public health issue that has a major impact on individuals, their families, workplaces and the community. The 2011-12 Australian Health Survey reports that 15.4% of Australians aged 15 years or older report are living with chronic pain. The prevalence of chronic pain is higher for women (16.9%) than it is for men (15.0%) and 68% of people with chronic pain are of working age. A recent report commissioned by Painaustralia estimated the cost of pain in Australia to be \$73.2 billion each year including health system costs, productivity losses and other costs (e.g. informal care, aids/modifications).

Overall objectives of the Chronic Pain Project¹

The overall objectives of the project are to:

1. Synthesise knowledge about the secondary prevention and management of chronic pain; and
2. Improve knowledge, knowledge-sharing and knowledge use among Primary Health Networks (PHNs) about options to address the secondary prevention and management of chronic pain in primary care.

Steering group

A small, time-limited project steering group involving lead clinicians, consumers, PHN and Local Health Network representatives and key researchers in the field was identified with Painaustralia (the peak national body for pain advocacy and policy) to provide rapid guidance and input across the course of the project. Steering group members are listed in **Appendix 1**.

Funding

The Chronic Pain Project at The Australian Prevention Partnership Centre is funded by the Medical Research Future Fund Boosting Preventive Health Research Program. Additional funding to support this project has been provided by the Sydney Medical School Foundation, University of Sydney.

Definitions

- **Pain** is defined as an unpleasant sensory and emotional experience associated with actual or potential tissue damage or described in terms of such damage.
- **Chronic pain** is defined as pain that lasts or recurs for more than three months. Chronic pain has recently been classified as a disease in itself by the World Health Organization, International Classification of Diseases (WHO-ICD-11).
- **Acute pain** is defined as pain that occurs immediately post-trauma or post-surgery. It is often self-limiting and usually resolves with healing within 3 months.
- **Subacute pain** is defined as the phase that lasts between six to twelve weeks post onset of acute pain.
- **Secondary prevention** of chronic pain is the early intervention of acute and subacute pain [herein referred to as (sub)acute] to prevent the progression to chronic pain and associated disability.

¹ The Australian Prevention Partnership Centre: Strategies and models for preventing or reducing the risk of the development of chronic pain in primary care (2018–2020)

Phase 1 of the Chronic Pain Project

In Phase 1 of the Chronic Pain Project, the research team conducted the following:

- i. Review of PHN recent Needs Assessments to assess whether chronic pain has been identified by PHNs as a health or service need and the key issues identified by PHNs related to chronic pain. For more information see the *Review of the Primary Health Networks Needs Assessments*.²
- ii. Consultation (interviews and a survey) with executive level staff and program managers from PHNs to understand the scope of work currently being implemented by PHNs related to the secondary prevention and management of chronic pain and to identify gaps. Twenty-five PHNs and one state PHN alliance (WA Primary Health Alliance³) participated in the consultation. For more information see the *Review of Primary Health Network Chronic Pain Initiatives: Summary of findings from the consultation with Primary Health Networks*.⁴

A framework of the types of chronic pain initiatives implemented in PHNs

In Phase 1 of the Chronic Pain Project, a framework of the different types of initiatives that PHNs were implementing related to the secondary prevention and management of chronic pain was developed. The framework is based on three goals adapted from the goals of the National Pain Strategy (Painaustralia)⁵ and aligned with PHNs' remit:

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

Goal 2: Ensuring health professionals are skilled and provide best-practice evidence-based care

Goal 3: Quality improvement and health system support

Results of the consultation with PHNs in Phase 1 of the project

The mapping of PHN chronic pain initiatives in Phase 1 of the project found a gap related to the secondary prevention of chronic pain with most initiatives currently being implemented by PHNs relating to the management of chronic pain. As a result of the gap highlighted in Phase 1, the opportunities for PHNs to improve the secondary prevention of chronic pain is the primary focus of Phase 2 of the Chronic Pain Project.

Purpose and scope of the rapid review

The purpose of the rapid review is to inform a deliberative dialogue with PHNs about the secondary prevention of chronic pain. A deliberative dialogue is an evidence-based method used to support policy making by discussing and contextualising research evidence in the light of the real-world experiences of policymakers.

Aims of the rapid review

The rapid review will seek to answer the following questions:

² De Morgan S, Blyth F and Walker P. Review of the Primary Health Networks Needs Assessments. The Australian Prevention Partnership Centre and the University of Sydney, September 2018.

³ WA Primary Health Alliance oversees the strategic commissioning functions of the three Western Australian Primary Health Networks: Perth North, Perth South and Country WA.

⁴ De Morgan S, Walker P and Blyth F. Review of Primary Health Network Chronic Pain Initiatives: Summary of findings from the consultation with Primary Health Networks. The Australian Prevention Partnership Centre and the University of Sydney, June 2019.

⁵ Pain Australia. National Pain Strategy 2010. Available at: www.painaustralia.org.au/the-national-pain-strategy/national-painstrategy.html

- i. What are the key principles related to the secondary prevention of chronic pain highlighted in the evidence?
- ii. What strategies related to Goal 1 (consumer and community initiatives) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)? Six key focus areas include:
 - a. pain that is multifocal or not condition specific
 - b. surgery
 - c. whiplash
 - d. low back pain
 - e. return to work and work-related injuries
 - f. opioid-related initiatives
- iii. What strategies related to Goal 2 (health professional capacity building) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)?
- iv. What strategies related to Goal 3 (health system support) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)?

The rapid review will include strategies implemented in the primary care setting as well as strategies implemented in hospital and compensable settings that have the potential to be adapted to the primary care setting.

Evidence sources published from 2010 to 2019 include the following:

- Peer review literature including clinical practice guidelines, systematic and narrative reviews, randomised controlled trials and protocols, and observational studies (Australia, UK, Europe, USA, Canada and New Zealand)
- Grey literature from key agencies in Australia and internationally
- Consultation with PHNs conducted in Phase 1 of the Chronic Pain Project
- Evidence identified by key stakeholders


The rapid review does not aim to systematically search for, or synthesise, all the relevant evidence related to the secondary prevention of chronic pain. The rapid review is a narrative synthesis of the evidence in the selected bodies of evidence and will highlight case study examples.

The evidence informs the development of the *Principles for the secondary prevention of chronic pain* and the mapping of options for PHNs to address the secondary prevention of chronic pain.

For more information about the search strategy see **Appendix 2**.

Principles for the secondary prevention of chronic pain

The principles for the secondary prevention of chronic pain were derived from a synthesis of relevant Australian and international guidelines, systematic reviews and narrative reviews. The *Principles for the secondary prevention of chronic pain* are outlined as follows:



See **Section 1** of the main report for the key elements of the *Principles for the secondary prevention of chronic pain*.

Prognostic risk factors for the progression from acute to chronic pain

The biopsychosocial model has been applied as a framework for understanding the complexity of progression of acute pain to chronic pain. Identifying people at risk of developing chronic pain is crucial. Risk factors for poor recovery are well documented and include psychological and social factors.

Risk factors can be identified early, and treatment can be tailored to the individual's risk profile to help prevent the progression to chronic pain. Screening tools and risk-based assessments can help to identify risk factors early.

Screening tools that have been used most widely include:

- The **STarT Back tool** has been used in primary care in the UK to identify low, medium or high risk of persistent disability in patients with low back pain and the Keele research group have developed online training to use the tool in routine care.
- The 10-item Swedish scale (the **Orebro Musculoskeletal Pain Screening Questionnaire - Short-Form - OMPSQ-SF**) has been used with injured workers in Australia with pain at any site, and training (online and face to face) is available for its use.
- A **clinical prediction rule** (CPR) has been used in the research setting in Australia to identify people who are at high risk of poor recovery in the early post-injury stage for people with whiplash-associated disorder.
- **Screening tools in the postoperative setting** are usually surgery-type specific and implemented in the hospital setting. One generic tool (a risk index) assesses the effect of 14 biomedical and psychosocial items that were derived from a systematic review of the CPSP risk factor literature.

However, there is a paucity of research related to the implementation of screening tools and risk-based interventions in the primary care setting. Education and training of primary care providers is needed so that clinicians understand the purpose of screening tools and other risk-based assessments within the context of their own clinical reasoning, how to communicate the results to patients and how the results can be used to inform the type of treatment needed for each patient.

For more information about risk factors, 'yellow flags' and screening tools see *Section 1: Principles for the secondary prevention of chronic pain* and for more information about the implementation of screening tools see *Section 2.2. Options related to Goal 2 (health professional capacity building)*.

Mapping of the options for Primary Health Networks

A range of options for the secondary prevention of chronic pain were developed based on the rapid review of the evidence and the consultations with PHNs conducted in Phase 1 of the project. The options have been grouped by the three goals of chronic pain initiatives implemented by PHNs, developed in Phase 1 of the project.

Each option will require different implementation considerations, organisational and behaviour change, funding and adaptation to the local PHN context. Key issues related to scalability, scale-up and sustainability will also need to be considered.

Mapping of the options for Primary Health Networks related to the secondary prevention of chronic pain

- Options relate to acute and subacute pain populations (e.g. post-surgery, post-injury, (sub)acute back pain) with a focus on people at risk of developing chronic pain.
- Options could be tailored to specific groups including Aboriginal and Torres Strait Islander peoples, people from culturally and linguistically diverse backgrounds, people from rural and remote areas, older Australians, people with dementia, children and young people and other relevant groups.

1. Options related to Goal 1 (consumer and community initiatives)

1.1 Face-to-face multidisciplinary consumer pain program (one or several group-based education sessions with/without individual consultation sessions with primary care providers)

The program may include:

- Group-based sessions (education and active, practice-based learning; physical activity; cognitive and behavioural strategies)
- Case management with individual consultation sessions with a primary care provider (e.g. physiotherapist, psychologist) as required
- Information resources (paper-based, links to online resources)

1.2 Psychologically-informed physical therapy program (individual consultation session(s) with a physiotherapist)

The program may include:

- Individual consultation sessions with a physiotherapist with psychologically-informed practice training (e.g. graded exercise and goal setting, cognitive and behavioural strategies, promotion of self-management)
- Information resources (paper-based, links to online resources)

1.3 Consumer initiative related to safe and effective use of medications; and non-initiation and tapering of opioids (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)

The initiative may include:

- Group based sessions (education and active, practice-based learning) may be embedded in a face-to-face multidisciplinary consumer pain program or a separate consumer workshop/education session(s)
- Or an online consumer initiative (e.g. webinar) including education and behavioural strategies
- Or individual consultation session(s) with a primary care provider e.g. GP, practice nurse, community pharmacist (education and behavioural strategies)

- Information resources (paper-based, links to online resources)

1.4 Transitions of care / pre-surgery consumer initiative (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)

The initiative may include:

- Individual sessions with a primary care provider e.g. GP, practice nurse. The sessions may include:
 - Pre-surgery risk assessment for developing chronic pain after surgery
 - Education about pain and pharmacological and non-pharmacological pain management
 - Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management
- Or an online consumer initiative (e.g. webinar) including education and behavioural strategies
- Information resources (paper-based, links to online resources)

1.5 Transitions of care / post-surgery consumer initiative (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)

The initiative may include:

- Individual sessions with a primary care provider e.g. GP, practice nurse, community pharmacist. The sessions may include:
 - Post-surgery risk assessment for developing chronic pain
 - A clinical medication review including medication reconciliation with active patient counselling
 - Education about pain and pharmacological and non-pharmacological pain management
 - Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management
 - Follow-up at, for example, 2 weeks, 6 weeks and 12 weeks
- Or an online consumer initiative (e.g. webinar) including education and behavioural strategies
- Information resources (paper-based, links to online resources)

1.6 Telehealth-assisted health care

- Delivery of healthcare at a distance using information communications technology (ICT).
- Telehealth connects clinicians or any other person(s) responsible for providing care to a patient and carer/s.
- Telehealth can be used for the purposes of assessment, intervention, consultation, education and/or supervision.
- Various models of care, for example,
 - Individual sessions connecting the patient (with/without their GP) to hospital-based clinician(s) (e.g. pain specialist, anaesthetist, surgeon, psychiatrist, nurse, physiotherapist)
 - Individual sessions connecting the patient with an allied health provider in the community such as a physiotherapist or clinical psychologist
 - Group sessions e.g. telerehabilitation (education, exercise)

1.7 Online consumer pain program (webinar or several online education sessions)

- The program is based on the same principles as face-to-face programs and uses online modules to teach pain management information and support patients to develop their self-management skills and encourage non-pharmacological approaches
- The program may have varying levels of telephone and/or email support from a primary care provider (e.g. psychologist, physiotherapist training in psychologically informed practice)

1.8 Mobile app

- Providing information and strategies similar to an online consumer pain program

1.9 Peer support group/network

- Delivered face-to-face, online or via social media

1.10 Community awareness campaign

- Delivered via social media, television, radio, print media or community events
- Campaign may aim to improve the community's understanding of pain; how to prevent chronic pain; promote self-management and non-pharmacological pain management; and promote safe and effective use of medicines.

1.11 Promotion of relevant consumer resources and programs implemented by other agencies

- E.g. paper-based and online information resources, online and face-to face consumer pain programs and support groups
- Promotion via consumer and health professional networks (events and newsletters), HealthPathways and online consumer distribution platforms (e.g. GoShare) and information portals (e.g. Patientinfo, Health Resource Directory)

2. Options related to Goal 2 (health professional capacity building)

2.1 Face-to-face and/or online education and training for GPs and other primary care providers

- Delivered face-to-face or online e.g. webinars, online modules or online platforms (Project ECHO)
- Education and training may include didactic sessions and case-based learning related to the following:
 - Explaining pain
 - Imaging
 - Risk factors for the progression of acute pain to chronic pain
 - Psychologically-informed practice (e.g. graded exercise and goal setting, cognitive and behavioural strategies, promotion of self-management)
 - Safe and effective use of medicines/opioid education
 - Follow-up of acute pain patients to monitor patient progress
 - Integration of care between hospital setting and primary care; and between primary care providers
 - Return to work

2.2 Opioid initiative about prescribing, non-initiation and deprescribing of opioids (face-to-face and/or online)

- Delivered face-to-face or online e.g. webinars, online modules or online platforms (Project ECHO)
- The initiative may include:
 - Didactic sessions and case-based learning
 - And/or behavioural strategies e.g. clinical reminder system

2.3 Interdisciplinary community of practice (CoP) (face-to-face and/or online)

- Delivered face-to-face or via online platform
- Chronic pain CoP or as part of a mental health CoP or Alcohol and Other Drugs (AOD) CoP
- And/or a network for primary care providers involved in a specific face-to-face consumer pain program

2.4 Promotion of relevant education and training and resources implemented by other agencies

- E.g. information resources, webinars, online modules, online platforms (Project ECHO), undergraduate and postgraduate training
- Promotion via health professional networks (events and newsletters) and HealthPathways

3. Options related to Goal 3 (health systems support initiatives)

3.1 Implementation of HealthPathways

- To assist general practitioners (GPs) with the management of patients with acute, subacute and chronic pain, and the referral of patients to specialists and allied health professionals

3.2 Transitions of care / health systems support initiative

The initiative may include:

- Electronic tools to facilitate quick, clear, and structured summary generation
- Use of electronic discharge notifications
- Online access to discharge information for general practitioners
- Discharge planning with shared involvement for follow-up of patients by hospital and community care providers

3.3 Prescription drug monitoring systems

- E.g. SafeScript is a real-time prescription monitoring and clinical decision support system that aims to provide doctors and pharmacists access to an up-to-the-minute medication supply history for certain high-risk medicines for their patient at the point of consultation

3.4 Electronic Persistent Pain Outcomes Collaboration (ePPOC)

- The Electronic Persistent Pain Outcomes Collaboration (ePPOC) is a national benchmarking system for the pain sector.
- It aims to improve clinical outcomes for people experiencing persistent pain through reporting and benchmarking.
- It is implemented and managed by the Australian Health Services Research Institute (AHSRI), University of Wollongong. It includes ePPOC for adults and PaedePPOC for children.
- ePPOC data is used to evaluate hospital-based pain services and community-based consumer pain programs

3.5 Other quality improvement systems

- E.g. The Practice Incentives Program (PIP), introduced in August 2019 by the Department of Health, Australian Government is a payment to general practices that participate in quality improvement activities to improve patient outcomes, and deliver best-practice care.

A narrative synthesis of the supporting evidence for each option related to each of the three goals is provided in the main body of the report. See **Appendix 3-5** for tables of the evidence informing each option.

Aboriginal and Torres Strait Islander people

The options above could be tailored to Aboriginal and Torres Strait Islander people. Suggestions for how to tailor the options to Aboriginal and Torres Strait Islander people based on the evidence include:

- Use a 'clinical yarning' approach for effective communication related to the assessment and management of pain. Clinical yarning is a patient-centred approach that marries Aboriginal cultural communication preferences with biomedical understandings of health and disease. Provide education and training for primary care providers about 'clinical yarning'. *For more information about clinical yarning see the main body of the report.*
- Ensure multidisciplinary group-based consumer pain programs involve education as well as behavioural skills approaches that focus on active, practice-based learning such as role play, feedback and small-group discussions; and are cofacilitated by an Aboriginal person (cultural knowledge) and primary care provider such as a physiotherapist (clinical knowledge), although an Aboriginal primary care provider could possess both these areas of knowledge.
- Develop culturally appropriate resources (e.g. booklets and videos about pain) in partnership with Aboriginal people and include, for example, Aboriginal people (and local Aboriginal people if possible) in the resources; visual formats; narratives; metaphors; and avoid medical jargon.
- Increase awareness among primary care providers of other potential barriers to access to care for Aboriginal and Torres Strait Islander people including institutional racism and discrimination; mistrust of health services; inadequate Aboriginal health staff; and financial barriers. Provide cultural safety training for primary care providers and support staff.

Resources for consumers and health professionals

- See **Section 4** in the main report for examples of consumer resources relevant to the secondary prevention of chronic pain.
- See **Section 5** in the main report for information and support to implement telehealth; and for examples of health professional education and training and other resources relevant to the secondary prevention of chronic pain.

Conclusion

The biopsychosocial model is widely accepted as the best approach to the assessment, secondary prevention and treatment of chronic pain.

Although risk factors for poor recovery are well documented and include psychological and social factors, there is a paucity of research about interventions to prevent chronic pain. There is also limited research about health professional capacity building and health system support initiatives related to the secondary prevention of chronic pain. Furthermore, very few initiatives related to the secondary prevention of chronic pain identified in this rapid review were implemented in the primary care setting.

However, some initiatives related to the secondary prevention of chronic pain implemented in the hospital and compensable settings have the potential to be adapted to the primary care setting. There are also many initiatives related to the management of chronic pain implemented by PHNs or reported in the peer-review and grey literature that have the potential to be adapted to the secondary prevention of chronic pain.

Overall, there are opportunities for PHNs as outlined in the mapping of the options for PHNs presented in this review to implement initiatives to improve the secondary prevention of chronic pain.

1. Rationale for the secondary prevention of chronic pain

Burden of disease of chronic pain

- Chronic pain is a substantial public health issue and chronic pain is increasing due to the ageing population.
- The 2011-12 Australian Health Survey reported that 15.4% of Australians aged 15 years or older report living with chronic pain. The prevalence of chronic pain is higher for women (16.9%) than it is for men (15.0%).
- Sixty-eight percent of people with chronic pain are of working age.⁶
- A recent report commissioned by Painaustralia estimated that 3.24 million Australians live with chronic pain and this number is projected to increase to 5.23 million by 2050.²
- The cost of pain in Australia is estimated to be \$73.2 billion each year including health system costs, productivity losses, other costs (e.g. informal care, aids/modifications).²
- Chronic pain for most sufferers has a major impact on individuals, their families, workplaces and the community.
- For the individual, chronic pain can lead to poorer quality of life, depression and anxiety, disability, loss of income and unemployment, impact on education, and feeling of stigmatisation and exclusion.

Aboriginal and Torres Strait Islander people

- Musculoskeletal pain is an important and poorly recognised issue in Aboriginal health care.(1)
- The prevalence of musculoskeletal pain conditions is higher amongst Aboriginal Australians when compared with non-Aboriginal Australians. The rate of back pain is 1.1 times higher; osteoarthritis is 1.2 to 1.5 times higher; and rheumatoid arthritis is 1.0 to 2.0 times higher.(1)
- Aboriginal people are potentially at higher risk of disabling musculoskeletal pain because disabling musculoskeletal pain conditions often co-exist with other health conditions and are associated with socioeconomic disadvantage.(1)

Focus areas: surgery, whiplash, low back pain, and return to work and work-related injuries

The burden of disease and the rationale for the secondary prevention of chronic pain is described below in the following focus areas: surgery, whiplash, low back pain, and return to work and work-related injuries.

Surgery

- The 11th revision of the International Classification of Diseases defines Chronic postsurgical pain as pain developing or increasing in intensity after a surgical procedure, in the area of the surgery, persisting beyond the healing process (that is, at least 3 months) and not better explained by another cause such as infection, malignancy, or a pre-existing pain condition.(2)

⁶ Deloitte Access Economics. The cost of pain in Australia. Painaustralia March 2018.

- Chronic postsurgical pain (CPSP) is common and may lead to significant disability.(3) The one year incidence of moderate to severe CPSP is approximately 12% for adults(4, 5) and approximately 22% for children.(4, 6)
- CPSP is a growing public health problem with 312 million major surgeries performed annually worldwide.(4, 7)

Whiplash

- Whiplash is a major health problem in Australia as well as other Western societies. It is the most common and costly injury following road traffic crashes with up to 50% of those injured still having pain or disability a year later.(8)
- Most recovery, if it occurs, takes place in the first 2-3 months postinjury, indicating that how people with whiplash are managed in the early stages will be critical to long-term outcome.(8, 9)
- Chronic whiplash affects peoples' quality of life, their family and social relationships, and impairs their ability to work. Consequently, whiplash has a huge economic impact.(10)

Low back pain

- A recent Lancet series (11-14) describes low back pain as a major global challenge requiring urgent action.
- Low back pain is an extremely common condition in populations worldwide and occurs in all age groups, from children to the elderly population. Most adults will have low back pain at some point.(14)
- Low back pain is now the number one cause of disability globally.(15) In 2015, the global point prevalence of activity-limiting low back pain was 7.3%, implying that 540 million people were affected at any one time. Disability from low back pain is highest in working age groups worldwide.
- The global burden of low back pain is projected to increase even further in coming decades, particularly in low-income and middle-income countries, straining health-care and social systems that are already overburdened.(14)
- Rarely can a specific cause of low back pain be identified; thus, most low back pain is termed non-specific.(14) Non-specific low back pain is estimated to be 90–95% of cases in primary care.
- Specific and serious causes of low back pain include vertebral fracture, inflammatory disorders such as axial spondyloarthritis, infection or malignancy.(14)
- Most episodes of low back pain improve substantially within 6 weeks, and by 12 months average pain levels are low.(16) However, approximately two-thirds of patients still report some pain at 3 months and 12 months.
- Recurrences of low back pain are common with approximately 33% of people having a recurrence within 1 year of recovering from a previous episode.(14)

Return to work and work-related injuries

- Soft tissue (musculoskeletal) injuries are the most common work-related injuries, estimated at 57.1% of all work-related injury/illness in Australia in 2018.(17)
- The vast majority (92.7%) of all workers surveyed in 2018 reported having returned to work at any time since their work-related injury or illness. However, one-in-five workers (19.6%) reported making more than one attempt to return to work, consequently having to take additional time off since returning to work, due to their work-related injury or illness).(17)
- Around three-in-eight (37.6%) workers who had returned to work reported that they worked reduced hours upon their return. Those who experienced mental illness were the most likely to work reduced hours upon returning to work (53.7%). Around three-in-eight (38.4%) workers who had returned to work

reported that they were performing slightly different/modified duties upon their return to work, while 19.0% reported performing completely different duties.(17)

- While little time is lost from work for most cases, a small proportion have delayed recovery and delayed return to work. If a worker is absent from work for 3 months or more following injury, the outlook becomes significantly more negative.(18) The longer an injured worker remains absent from work the greater is their risk of never returning to work; longer term ill-health and financial insecurity; and costs to the community.(19, 20)
- In general, work is good for long-term health and well-being.(18, 21)

Opioids

The rationale for non-initiation or tapering of opioids in the shortest period possible post-surgery, post-injury and for (sub)acute low back pain:

- Opioids are not recommended for chronic pain and have led to harm such as addiction, overdose and death.
- Current practices for the prescription of opioids at discharge after surgery are highly variable and often excessive, elevating the risk of opioid dependence.(22)
- Many surgical procedures are associated with increased risk of chronic opioid use (in people that have not used an opioid in the year prior to surgery).(23)
- High postsurgical opioid consumption is also a risk factor for chronic postsurgical pain.(4, 24)
- *Acute Pain Management: Scientific Evidence (2015)* by the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine highlights that short-term opioid therapy may lead to long-term opioid use (3) and suggests a “universal precaution” approach for opioid prescribing after surgery in the setting of prescribing discharge medications.
- *US Clinical Practice Guidelines for Pain Management in Acute Musculoskeletal Injury (2019)* recommend prescribing the lowest effective immediate-release opioid for the shortest period possible and connecting patients to psychosocial interventions as indicated.(25) The guidelines also recommend opioid education for prescribers and patients.
- After surgery and injury, short-term opioid therapy may lead to long-term opioid use.
- Continued use of opioids after work-related injuries delays recovery and poorer return to work outcomes.

2. Principles for the secondary prevention of chronic pain

The *Principles for the secondary prevention of chronic pain* were derived from a synthesis of relevant Australian and international guidelines(3, 25-29) and systematic and narrative reviews.(2, 4, 13, 14, 18, 22, 30-41) The *Principles for the secondary prevention of chronic pain* are outlined as follows:




Figure 1: Principles for the secondary prevention of chronic pain

Key elements

See **Figure 2** for the key elements of the *Principles for the secondary prevention of chronic pain*.

Principles for the secondary prevention of chronic pain

Principle 1: Biopsychosocial model of pain

- Pain is highly personal and subjective experience
- **Biomedical**- what's happening in your body
- **Psycho**- what's happening to you as a person
- **Socio**- what's happening in your world

Principle 2: Recognition of prognostic risk factors

- **Recognition of risk factors** for the progression of (sub)acute pain to chronic pain including demographic, lifestyle, genetic, clinical, surgery related and psychological factors
- **Surgery:** Preoperative, perioperative and transitional phase patient assessment and screening (consider using a surgery-type specific screening tool or generic tool)
- **Post-injury and (sub)acute back pain:** Patient assessment on the first contact with primary care provider and screening (consider using a screening tool (e.g. Orebro, STarT Back,))

Principles 3: Multidisciplinary and multimodal approach tailored to the individual

- **Tailoring treatment** to individual's risk profile
- **Encouraging self-management and non-pharmacological options**
- **Multidisciplinary and multimodal** including:
 - Patient education and reassurance
 - Movement/staying active and pacing
 - Physical therapies (e.g. manual therapy, exercise programs)
 - Mind-based therapies to understand the relationship between beliefs and behaviours and develop goal-orientated plans (e.g. cognitive behavioural therapy, psychologically informed physiotherapy, stress-management)
 - Pharmacological: safe and effective use of medications; opioid education for consumers including tapering; and use of simple analgesic medicines
- **Surgery:** preoperative and perioperative education about pain management and behavioural strategies including coping methods and goal setting for postoperative phase
- **Increase access to care by telehealth and digitally delivered treatments**

Principles 6: Public awareness

- **Explaining pain**
- **Risk factors for progression** of (sub)acute pain to chronic pain
- **Promoting self-management and non-pharmacological approaches**
- **Safe and effective use of medicines** (e.g. opioid education)

Principle 5: Improved coordination of care between the acute setting and primary care and between primary care providers

- Greater communication **between hospital services and general practice** (e.g. discharge/transitional phase planning)
- Greater communication between **primary care providers** (e.g. GPs and allied health practitioners, practice nurses, community pharmacists)

Principles 4: Education and training of primary care providers (GPs, practice nurses, allied health practitioners, community pharmacists)

- **Explaining pain:** How to explain pain to patients and language to avoid/use
- **Imaging:** When to use imaging and how to explain imaging results to patients
- **Risk factors for the progression of acute pain to chronic pain:**
 - How to assess patients for risk factors for poor recovery to inform type of treatment ('yellow flags')
 - How to use specific screening tools in the context of clinician's own clinical reasoning
- **Psychologically informed practice:**
 - How to help patients set goals and promote self-management, pacing and other non-pharmacological approaches
 - How to apply simple coping methods and behavioural strategies
 - Referring to allied health practitioners (e.g. physiotherapists, psychologists)
 - Additional training for physiotherapists- psychologically informed physical therapy
- **Safe and effective use of medicines/opioid education:**
 - Appropriate opioid prescribing
 - Non-initiation and deprescribing of opioids
 - Non-opioid pain analgesic medicines
- **Follow-up of acute pain patients to monitor patient progress:** Scheduling times for follow-up (e.g. 2, 6 and 12 wks)
- **Integration of care between hospital setting and primary care; and between primary care providers**
 - Communicating with hospital services and other primary care providers
 - Timely referral to specialist services (e.g. at 12 weeks) if appropriate
- **Return to work:**
 - Approach to patients
 - Capacity certification
 - Understanding the compensable environment

Figure 2: Key elements of the principles for the secondary prevention of chronic

Recognition of prognostic risk factors

The biopsychosocial model has been applied as a framework for understanding the complexity of progression of acute pain to chronic pain. Identifying people at risk of developing chronic pain is crucial. Risk factors for poor recovery are well documented and include psychological and social factors. Many risk factors are modifiable.

Risk factors can be identified early, and treatment can be tailored to the individual's risk profile to help prevent the progression to chronic pain.(41)

Chronic postsurgical pain

Chronic postsurgical pain (CPSP) is defined as pain developing or increasing in intensity after a surgical procedure, in the area of the surgery, persisting beyond the healing process (i.e. at least 3 months) and not better explained by another cause such as infection, malignancy, or a pre-existing pain condition (World Health Organization, International Classification of Diseases WHO-ICD-11).

A recent Lancet series describes postoperative pain management and risk factors for transition from acute to chronic pain after surgery.(2) Risk factors for the development of CPSP are well documented (2-4, 42, 43) and include the severity of presurgical chronic pain and postsurgical acute pain, intraoperative nerve injury and psychological factors such as anxiety, pain catastrophising, depression, psychological vulnerability and stress.

Five core risk factor domains have been identified by the Initiative on Methods, Measurement, and Pain Assessment in Clinical Trials (IMMPACT): demographic, genetic, clinical, surgery related, and psychological.(44, 45) Risk factors for CPSP are not independent of each other, but interlinked and include the following:

1. Demographics and lifestyle

- Age
- Gender
- Marital status or living arrangements
- Education level
- Employment status
- Compensation status
- Obesity
- Smoking

2. Genetic

- Candidate gene mutations associated with increased pain

3. Clinical

- Surgical factors, including type of surgery, surgical technique (open vs laparoscopic), duration of surgery, type of anaesthesia (general vs regional), and perioperative
- Analgesic regimen (systemic vs spinal and pre-emptive); surgical complications and re-operating
- Medical comorbidities
- Previous disability or pain interference

4. Preoperative pain (area of operation or elsewhere)

5. Postoperative pain (intensity and duration)

6. Psychological

- Fear or anxiety
- Depression
- Pain catastrophising
- Other psychological issues (e.g. vulnerability factors)

Figure 3 outlines the risk factors for the development of chronic postsurgical pain.

Screening tools are usually surgery-type specific.(2) One generic tool (risk index) assesses the effect of 14 biomedical and psychosocial items that were derived from a systematic review of the CPSP risk factor literature.(2, 46)



Figure 3: Risk factors for the development of chronic postsurgical pain

Low back pain

A recent Lancet series (11-14) describes low back pain as a major global challenge requiring urgent action. The series describes the epidemiology of low back pain, the complexity of the condition, risk factors for progression of acute to chronic pain and the evidence to support interventions to improve the prevention and management of low back pain.

The biopsychosocial model has been applied as a framework for understanding the complexity of low back pain disability. Low back pain is a complex condition with multiple contributors to both the pain and associated disability, including psychological factors (e.g. depression, catastrophising, fear avoidance beliefs), social factors (e.g. physical work-loads, education, compensation, work satisfaction), biophysical factors (e.g. previous episodes, back pain intensity and presence of leg pain), comorbidities, and pain-processing mechanisms. Lifestyle factors, such as smoking, obesity, and low levels of physical activity, that relate to poorer general health, are also associated with occurrence of low back pain episodes.(14)

Figure 4 outlines the risk factors for low back pain and associated disability.




Figure 4: Risk factors for low back pain and associated disability

The NSW Agency for Clinical Innovation (ACI) has recently developed a model of care (MOC) for acute low back pain (2016) that can be applied to primary care and emergency department settings.(28) The MOC supports the identification of ‘yellow flags’ (41) (see **Figure 5**) to guide the level and type of treatment recommendation and prevent the progression to chronic pain. The MOC recommends using screening tools such as:

- The STarT Back tool (47, 48) has been used in primary care in the UK to identify low, medium or high risk of persistent disability in patients with low back pain and the Keele group have developed online training to use the tool in routine care.
- The 10-item Swedish scale (the Orebro Musculoskeletal Pain Screening Questionnaire – Short-Form – OMPSQ-SF) (19, 20) has been used in injured workers in Australia with pain at any site. Training (online and face to face) is available for its use.

<p>Yellow flags: assessed using either the STarT Back or Örebro tools</p> 	<ul style="list-style-type: none"> ▶ Belief that pain and activity are harmful ▶ “Sickness behaviours” (like extended rest) ▶ Low or negative moods, social withdrawal ▶ Treatment that does not fit with best practice ▶ Problems with compensation system ▶ Previous history of back pain, time off work, other claims ▶ Problems at work, poor job satisfaction ▶ Overprotective family or lack of social support. 	<p>Further information concerning the Yellow Flags can be found in the <i>New Zealand acute low back pain guideline</i> available at: http://www.acc.co.nz/PRD_EXT_CSMP/groups/external_communications/documents/guide/prd_ctrb112930.pdf</p>
--	---	---

Figure 5: Yellow flags

Whiplash-associated disorder

Current research for acute whiplash-associated disorder is investigating a risk stratification approach to care similar to the approach used in low back pain. The physiotherapist (the most commonly used practitioner delivering care to patients with acute whiplash-associated disorder) provides care that addresses the physical and psychological factors identified in a detailed assessment of the patient.

A clinical prediction rule (CPR) has been validated which can identify patients from the early acute post injury stage who are at high risk of poor recovery at 12 months. The tool can also identify those who are likely to fully recover and a third medium risk group who could either recover or develop chronic pain and disability.(49) However, a qualitative study evaluating the perceptions of physiotherapists, chiropractors and osteopaths of adopting the clinical prediction rule found that clinicians’ understanding and use of the tool was mixed. The authors suggest that further education is needed so that clinicians understand the purpose of the tool, how to use the tool in the context of their own clinical reasoning and how to communicate the results to patients.(50)

Return to work and work-related injuries

Psychological and social/environmental factors are predictors for delayed recovery and disability associated with chronic pain for injured workers.(19) Legal involvement in an injury and an associated compensation claim following an injury are also associated with worse physical and psychological functional outcomes.(51)

A validation study of Örebro Musculoskeletal Pain Screening Questionnaire-short version (ÖMPSQ-SF) to predict time to return to pre-injury work duties following a work-related soft tissue injury (regardless of body location) provides strong support for the use of the ÖMPSQ-SF in an applied setting for identifying those injured workers likely to have delayed return to work when administered within 15 days of the injury.(20)

Overall comments

Identifying people at risk of developing chronic pain is crucial. Risk factors for poor recovery are well documented and include psychological and social factors. Risk factors can be identified early, and treatment can be tailored to the individual’s risk profile to help prevent the progression to chronic pain. However, there is a paucity of research related to the implementation of screening tools and risk-based interventions in the primary care setting.

Education and training of primary care providers is needed so that clinicians understand the purpose of screening tools and other risk-based assessments within the context of their own clinical reasoning, how to communicate the results to patients and how the results can be used to inform the type of treatment needed for each patient.

3. Mapping of the options for Primary Health Networks with supporting evidence

A range of options for the secondary prevention of chronic pain were developed based on the rapid review of the evidence and the consultations with PHNs conducted in Phase 1 of the project. The options have been stratified by the three goals of chronic pain initiatives implemented by PHNs, developed in Phase 1 of the project.

Each option will require different implementation considerations, organisational and behaviour change, funding and adaptation to the local PHN context.

Table 1 maps the options for PHNs related to the secondary prevention of chronic pain.

Supporting evidence

Although risk factors for poor recovery are well documented and include psychological and social factors, there is a paucity of research about interventions to prevent chronic pain. There is also limited research about health professional capacity building and health system support initiatives related to the secondary prevention of chronic pain. Furthermore, very few initiatives related to the **secondary prevention of chronic pain** identified in this rapid review were implemented in the **primary care setting**.

However, some initiatives related to the secondary prevention of chronic pain implemented in the **hospital and compensable settings** have the potential to be adapted to the primary care setting. There are also many initiatives related to the **management of chronic pain** implemented by PHNs or reported in the peer-review and grey literature that have the potential to be adapted to the secondary prevention of chronic pain.

A narrative synthesis of the supporting evidence for each option in each of the three goals is provided in this section of the report. See **Appendix 3-5** for tables of the evidence informing each option.

Evidence sources include:

- Peer review literature including clinical practice guidelines, systematic and narrative reviews, randomised controlled trials and protocols, and observational studies (Australia, UK, Europe, USA, Canada and New Zealand)
- Grey literature from key agencies in Australia and internationally
- Consultation with PHNs conducted in Phase 1 of the Chronic Pain Project
- Evidence identified by key stakeholders

The rapid review does not aim to systematically search for, or synthesise, all the relevant evidence related to the secondary prevention of chronic pain. *For more information about the search strategy see **Appendix 2**.*



Table 1: Map of the options for PHNs related to the secondary prevention of chronic pain

<ul style="list-style-type: none"> Options relate to acute and subacute pain populations (e.g. post-surgery, post-injury, (sub)acute back pain) with a focus on people at risk of developing chronic pain. Options could be tailored to specific groups including Aboriginal and Torres Strait Islander peoples, people from culturally and linguistically diverse backgrounds, people from rural and remote areas, older Australians, people with dementia, children and young people and other relevant groups.
<p>1. Options related to Goal 1 (consumer and community initiatives)</p>
<p>1.1. Face-to-face multidisciplinary consumer pain program (one or several group-based education sessions with/without individual consultation sessions with primary care providers)</p> <p>The program may include:</p> <ul style="list-style-type: none"> Group-based sessions (education and active, practice-based learning; physical activity; cognitive and behavioural strategies) Case management with individual consultation sessions with a primary care provider (e.g. physiotherapist, psychologist) as required Information resources (paper-based, links to online resources)
<p>1.2 Psychologically-informed physical therapy program (individual consultation session(s) with a physiotherapist)</p> <p>The program may include:</p> <ul style="list-style-type: none"> Individual consultation sessions with a physiotherapist with psychologically-informed practice training (e.g. graded exercise and goal setting, cognitive and behavioural strategies, promotion of self-management) Information resources (paper-based, links to online resources)
<p>1.3 Consumer initiative related to safe and effective use of medications; and non-initiation and tapering of opioids (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)</p> <p>The initiative may include:</p> <ul style="list-style-type: none"> Group based sessions (education and active, practice-based learning) may be embedded in a face-to-face multidisciplinary consumer pain program or a separate consumer workshop/education session(s) Or an online consumer initiative (e.g. webinar) including education and behavioural strategies Or individual consultation session(s) with a primary care provider e.g. GP, practice nurse, community pharmacist (education and behavioural strategies) Information resources (paper-based, links to online resources)
<p>1.4 Transitions of care / pre-surgery consumer initiative (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)</p> <p>The initiative may include:</p> <ul style="list-style-type: none"> Individual consultation sessions with a primary care provider e.g. GP, practice nurse. The sessions may include: <ul style="list-style-type: none"> Pre-surgery risk assessment for developing chronic pain after surgery Education about pain and pharmacological and non-pharmacological pain management Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management Or an online consumer initiative (e.g. webinar) including education and behavioural strategies Information resources (paper-based, links to online resources)
<p>1.5 Transitions of care / post-surgery consumer initiative (a group-based education session or webinar; and/or individual consultation session(s) with a primary care provider)</p> <p>The initiative may include:</p>

- Individual consultation sessions with a primary care provider e.g. GP, practice nurse, community pharmacist. The sessions may include:
 - Post-surgery risk assessment for developing chronic pain
 - A clinical medication review including medication reconciliation with active patient counselling
 - Education about pain and pharmacological and non-pharmacological pain management
 - Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management
 - Follow-up at, for example, 2 weeks, 6 weeks and 12 weeks
- Or an online consumer initiative (e.g. webinar) including education and behavioural strategies
- Information resources (paper-based, links to online resources)

1.6 Telehealth-assisted health care

- Delivery of healthcare at a distance using information communications technology (ICT).
- Telehealth connects clinicians or any other person(s) responsible for providing care to a patient and carer/s.
- Telehealth can be used for the purposes of assessment, intervention, consultation, education and/or supervision.
- Various models of care, for example,
 - Individual sessions connecting the patient (with/without their GP) to hospital-based clinician(s) (e.g. pain specialist, anaesthetist, surgeon, psychiatrist, nurse, physiotherapist)
 - Individual sessions connecting the patient with an allied health provider in the community such as a physiotherapist or clinical psychologist
 - Group sessions e.g. telerehabilitation (education, exercise)

1.7 Online consumer pain program (a webinar or several online education sessions)

- The program is based on the same principles as face-to-face programs and uses online modules to teach pain management information and support patients to develop their self-management skills and encourage non-pharmacological approaches
- The program may have varying levels of telephone and/or email support from a primary care provider (e.g. psychologist, physiotherapist training in psychologically informed practice)

1.8 Mobile app

- Providing information and strategies similar to an online consumer pain program

1.9 Peer support group/network

- Delivered face-to-face, online or via social media

1.10 Community awareness campaign

- Delivered via social media, television, radio, print media or community events
- Campaign may aim to improve the community's understanding of pain; how to prevent chronic pain; promote self-management and non-pharmacological pain management; and promote safe and effective use of medicines.

1.11 Promotion of relevant consumer resources and programs implemented by other agencies

- E.g. paper-based and online information resources, online and face-to face consumer pain programs and support groups
- Promotion via consumer and health professional networks (events and newsletters), HealthPathways and online consumer distribution platforms (e.g. GoShare) and information portals (e.g. Patientinfo, Health Resource Directory)

2. Options related to Goal 2 (health professional capacity building)

2.1 Face-to-face and/or online education and training for GPs and other primary care providers

- Delivered face-to-face or online e.g. webinars, online modules or online platforms (Project ECHO)

- Education and training may include didactic sessions and case-based learning related to the following:
 - Explaining pain
 - Imaging
 - Risk factors for the progression of acute pain to chronic pain
 - Psychologically-informed practice (e.g. graded exercise and goal setting, cognitive and behavioural strategies, promotion of self-management)
 - Safe and effective use of medicines/opioid education
 - Follow-up of acute pain patients to monitor patient progress
 - Integration of care between hospital setting and primary care; and between primary care providers
 - Return to work

2.2 Opioid initiative about prescribing, non-initiation and deprescribing of opioids (face-to-face and/or online)

- Delivered face-to-face or online e.g. webinars, online modules or online platforms (Project ECHO)
- The initiative may include:
 - Didactic sessions and case-based learning
 - And/or behavioural strategies e.g. clinical reminder system

2.3 Interdisciplinary community of practice (CoP) (face-to-face and/or online)

- Delivered face-to-face or via online platform
- Chronic pain CoP or as part of a mental health CoP or Alcohol and Other Drugs (AOD) CoP
- And/or a network for primary care providers involved in a specific face-to-face consumer pain program

2.4 Promotion of relevant education and training and resources implemented by other agencies

- E.g. information resources, webinars, online modules, online platforms (Project ECHO), undergraduate and postgraduate training
- Promotion via health professional networks (events and newsletters) and HealthPathways

3. Options related to Goal 3 (health systems support initiatives)

3.1 Implementation of HealthPathways

- To assist general practitioners (GPs) with the management of patients with acute, subacute and chronic pain, and the referral of patients to specialists and allied health professionals

3.2 Transitions of care / health systems support initiative

The initiative may include:

- Electronic tools to facilitate quick, clear, and structured summary generation
- Use of electronic discharge notifications
- Online access to discharge information for general practitioners
- Discharge planning with shared involvement for follow-up of patients by hospital and community care providers

3.3 Prescription drug monitoring systems

- E.g. SafeScript is a real-time prescription monitoring and clinical decision support system that aims to provide doctors and pharmacists access to an up-to-the-minute medication supply history for certain high-risk medicines for their patient at the point of consultation

3.4 Electronic Persistent Pain Outcomes Collaboration (ePPOC)


- The Electronic Persistent Pain Outcomes Collaboration (ePPOC) is a national benchmarking system for the pain sector.
- It aims to improve clinical outcomes for people experiencing persistent pain through reporting and benchmarking.

- It is implemented and managed by the Australian Health Services Research Institute (AHSRI), University of Wollongong. It includes ePPOC for adults and PaedePPOC for children.
- ePPOC data is used to evaluate hospital-based pain services and community-based consumer pain programs

3.5 Other quality improvement systems

- E.g. The Practice Incentives Program (PIP), introduced in August 2019 by the Department of Health, Australian Government is a payment to general practices that participate in quality improvement activities to improve patient outcomes, and deliver best-practice care.

3.1 Options related to Goal 1 (consumer and community initiatives)



3.1.1 Face-to-face multidisciplinary consumer pain program

- A face-to-face multidisciplinary consumer pain program could be implemented by PHNs for consumers with (sub)acute pain at risk of developing chronic pain with referrals by GP, specialist or allied health practitioners (with GP final sign-off)
 - Multidisciplinary treatments are treatments that target physical as well as psychological or social aspects of pain and involve a team of healthcare providers with different professional backgrounds and training (Martin et al 2017 Cochrane Database of Systematic Reviews)
- Consumer pain programs usually include:
 - Group based sessions (education and active, practice-based learning; physical activity; cognitive and behavioural strategies); and
 - Case management and individual sessions with primary care providers (e.g. physiotherapist, psychologist) as required; and
 - Information resources (paper-based, links to online resources)
- A consumer pain program could also be tailored to:
 - Specific (sub)acute consumer pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]
 - Specific groups including Aboriginal and Torres Strait Islander peoples, people from culturally and linguistically diverse backgrounds, people from rural and remote areas, older Australians, people with dementia, children and young people and other relevant groups. For example,
 - Ensure multidisciplinary group-based consumer pain programs for Aboriginal and Torres Strait Islander people are cofacilitated by an Aboriginal person (cultural knowledge) and


primary care provider such as a physiotherapist (clinical knowledge), although an Aboriginal primary care provider could possess both these areas of knowledge.

- Develop culturally appropriate resources (e.g. booklets and videos about pain) for Aboriginal and Torres Strait Islander people in partnership with Aboriginal people and include, for example, Aboriginal people (and local Aboriginal people if possible) in the resources; visual formats; narratives; metaphors; and avoid medical jargon.

*For a summary of the enablers to implementing consumer pain programs identified in the consultation with PHNs in Phase 1 of the Chronic Pain Project see **Appendix 6**.*

Supporting evidence

Face-to-face multidisciplinary consumer pain program



3.1.2 Psychologically informed physical therapy program

- A psychologically informed physical therapy individual program could be implemented by PHNs for consumers with (sub)acute pain at risk of developing chronic pain with referrals by GP or specialist to a physiotherapist.
- A psychologically informed physical therapy individual program usually includes:
 - Individual sessions with a physiotherapist with psychologically informed practice training (e.g. graded exercise and goal setting, cognitive and behavioural strategies, promotion of self-management); and
 - Information resources (paper-based, links to online resources)
- PHNs could provide additional psychologically informed physical therapy training for physiotherapists e.g. three-step training approach: a treatment manual, an experiential workshop, and ongoing supervision with consultation and feedback; or promote education and training offered by other agencies.

- Initiative could be tailored to specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence

Psychologically informed physical therapy program



Peer review literature

- Brunner 2013 systematic review- cognitive behaviour therapy-based treatments - (sub)acute low back pain
- Hall 2018 systematic review - physiotherapist-led cognitive-behavioural interventions - low back pain
- Nicholls 2018 systematic review – perioperative cognitive behavioural therapy
- Foster 2014 (IMPACT Back) ---stratified care-psychologically informed physical therapy - (sub)acute and chronic pain
- Sterling 2019 (StressModex) - physiotherapist-led intervention of stress inoculation training and exercise - acute whiplash
- Sullivan 2006 -psychological intervention and physical therapy in whiplash
- Archer 2016 - cognitive-behavioural based physical therapy – post-surgery
- Nicholas 2019 -early intervention for injured workers- compensation environment

Narrative synthesis

Focus area 1: multifocal or not condition specific

- The **biopsychosocial model** is widely accepted as the best approach to the assessment, prevention and treatment of chronic pain.(52, 53)

Management

- Although risk factors can identify individuals that are likely to develop chronic pain, there is a **lack of research about interventions to prevent chronic pain**.(44)
- **Face-to-face multidisciplinary consumer pain programs** that incorporate physical and psychological strategies are effective for people with chronic pain.(54, 55) However, there is limited research about face-to-face multidisciplinary consumer pain programs for people with (sub)acute pain (not-condition specific) at risk of developing chronic pain. A potential prototype has been developed for adapting the *Turning Pain Into Gain* consumer pain program for consumers with (sub)acute pain at risk of developing chronic pain. *For more information see Appendix 9.*
- A multidisciplinary pain management service for people at risk of developing chronic pain after surgery (as part of a **transitional pain service in Canada**) has been shown to be effective. *See below Focus area 2: Surgery.*
- There has been growing interest in **psychologically oriented pain management** over the past three to four decades.(56) Primary care physicians and physical therapists have delivered psychologically informed

practice as part of a stratified care approach involving screening and targeting of treatment for people at high risk for pain-associated disability. See below *Focus area 4: (Sub)acute low back pain*.

- A recent systemic review(36) about the secondary prevention of chronic musculoskeletal pain identified nine studies. Most studies included patients with (sub)acute low back pain. Most interventions included physical and psychological strategies provided by physiotherapists with mind-based training. **Stratified programs** showed significant improvements compared with a “one-size-fits-all” treatment in several domains of the International Classification of Functioning, Disability and Health. The authors concluded that simple educational messages seemed sufficient for low-risk patients; medium- and high-risk patients benefited from a physical reactivation programme combined with education; and in high-risk patients, an additional cognitive-behavioural intervention further improved the outcome.
- **Pain neuroscience education** has been found, in a recent systematic review (40), to facilitate the ability of chronic musculoskeletal pain patients to cope with their condition but it does not produce clinically significant decreases in pain, disability, kinesiophobia or catastrophizing. A recent clinical trial also found that intensive patient education (information on pain and biopsychosocial contributors plus self-management techniques, such as remaining active and pacing) in patients receiving first-line care for acute back pain (advice, reassurance, and simple analgesia, if necessary) may be no better than active listening in improving pain outcomes.(57)

Relevant research currently being conducted

- (Australia) Implementation of a clinical PATHway of CarE to improve patient health outcomes and reduce costs for common musculoskeletal disorders (low back pain, neck pain or whiplash, knee osteoarthritis) in primary care (**PACE study**). The PACE intervention aims to identify patients at risk of poor prognosis to improve their management in primary health care settings using a stratified care model. Patients are identified within four weeks of seeking care. The 10-item Orebro Musculoskeletal pain questionnaire will be used as the generic measure to assess risk of poor prognosis for all conditions. Validated condition-specific tools where available (e.g. STarT Back for low back pain; prognostic prediction tool for whiplash) will be used to cross reference and ensure appropriate stratification. The team is also currently working on an additional prognostic screening tool for all musculoskeletal conditions that may also be used for cross validation.

Specific population groups

- The National Strategic Action Plan for Pain Management developed by PainAustralia(58) recommends that prevention and early intervention strategies include initiatives **relevant to specific groups** including Aboriginal and Torres Strait Islander peoples, people from culturally and linguistically diverse backgrounds, people from rural and remote areas, older Australians, people with dementia, children and young people and other relevant groups.

Aboriginal and Torres Strait Islander people

- A recent editorial highlights that **musculoskeletal pain is an important and poorly recognised issue in Aboriginal health care**.(1) Despite this, **access to care is low**. For example, Aboriginal people with hip and knee osteoarthritis access primary care at around half the rate of non-Aboriginal people (3.2 versus 6.5 per 1000 encounters for knee osteoarthritis, 1.2 versus 2.3 per 1000 encounters for hip osteoarthritis) and rates are lower again for hip/knee arthroplasty.(1) Although there are limited data for other musculoskeletal pain conditions or physiotherapy, the authors suggest that access to care is likely to be similarly low. Yet interventions provided by physiotherapist, such as facilitating self-management, exercise and activity, could help prevent and better manage chronic pain and other chronic conditions.
- **‘Clinical yarning’** can be used as a framework to improve communication with Aboriginal people about musculoskeletal pain.(1, 59) There is cultural and research evidence that supports this approach. Clinical yarning has the potential to improve outcomes for patients and practitioners.

- Clinical yarning is a patient-centred approach that marries Aboriginal cultural communication preferences with biomedical understandings of health and disease. Clinical yarning consists of three interrelated areas: the social yarn, in which the practitioner aims to find common ground and develop the interpersonal relationship; the diagnostic yarn, in which the practitioner facilitates the patient's health story while interpreting it through a biomedical or scientific lens; and the management yarn, that employs stories and metaphors as tools for patients to help them understand a health issue so a collaborative management approach can be adopted.

See Focus area 4: (sub)acute back pain below for information about an intervention with Aboriginal people with back pain.

Focus area 2 - surgery

Management

- **Optimal management of postoperative pain begins in the preoperative period.** *For more information about education and behavioural strategies in the preoperative phase to reduce chronic postsurgical pain see Section 2.3.2 Transitional care/discharge planning initiative.*
- **Early postoperative exercise training and rehabilitation** is recommended for better functional outcomes and enhanced recovery.(33, 60)
- **There is a potential role for a transitional pain clinic** which aims to overcome the disconnect between ward-based acute postoperative pain management and outpatient chronic pain management with a comprehensive and integrated pain service that identifies patients at risk of chronic pain.(2)
 - The Toronto General Hospital Transitional Pain Service (TPS) is an example of this model. Patients are identified early through screening for physical and mental health problems e.g. anxiety, depression, pain catastrophizing, chronic opioid use, pre-existing chronic pain, severe postsurgical pain, high postsurgical opioid consumption. Patients are provided comprehensive care by a multidisciplinary team consisting of pain physicians, advanced practice nurses, psychologists, and physical therapists. Clinical services at the TPS include multimodal medication optimisation by anaesthesiologists, postsurgical physical therapy and acupuncture, and a pain psychology intervention consisting of pain education, mindfulness training, brief hypnosis, and a form of cognitive behavioural treatment called acceptance and commitment therapy (ACT). Preliminary evidence indicates that the TPS effectively reduces pain intensity, pain-related interference, pain catastrophising, symptoms of anxiety and depression, and opioid use.(4, 24)

Focus area 3 – (sub)acute whiplash

Management

- Clinical guidelines for the management of acute whiplash-associated disorders(27) recommend exercise and activity, but systematic reviews have concluded that exercise/activity alone provides only a small benefit.(10, 61)
- Current research for acute whiplash-associated disorder is investigating **a risk stratification approach to care** similar to the approach used in low back pain. The physiotherapist (the most commonly used practitioner delivering care to patients with acute whiplash-associated disorder) provides care that addresses the physical and psychological factors identified in a detailed assessment of the patient.
- An example of this approach is a recent trial (StressModex) of a **physiotherapist-led intervention of stress inoculation training** (teaching strategies to assist participants in managing acute stress responses) **and exercise** for patients with acute whiplash-associated disorder at risk of poor recovery (moderate pain-related disability and hyperarousal symptoms).(10) The intervention resulted in clinically relevant

improvements in pain-related disability, stress, depressive symptoms, pain self-efficacy and perceived recovery compared with exercise alone, the most commonly recommended treatment for acute WAD. The treatment benefit was maintained at 12-month follow-up. The authors conclude that physiotherapists with some additional training, could apply stress inoculation training for whiplash-injured patients at risk of poor recovery. See 3.2.1 Face-to-face and/or online education and training for more information about the training of physiotherapists to implement stress-inoculation.

For information about screening tools see 3.2.1 Face-to-face and/or online education and training.

Research currently being conducted

- An Australian multi-centre, randomised controlled trial (**Whiplash ImPaCT**) involves people within six weeks of their whiplash injury and their primary care providers (including general practitioners, physiotherapists, chiropractors, or osteopaths).(9) Participants are screened for risk of poor recovery using the clinical prediction rule for ongoing pain and disability and provided with care to match their predicted risk. Participants at low risk of ongoing pain and disability (hence, predicted to fully recover) will receive up to three sessions of guideline-based advice and exercise with their primary healthcare provider. Participants at medium/high risk of developing ongoing pain and disability will be referred to a specialist (defined as a practitioner with expertise in whiplash) who will conduct a more in-depth physical and psychological assessment. As a result, the specialist will liaise with the original primary healthcare provider and determine one of three further pathways of care: 1) continue current care; 2) exercise-based physical therapy with psychologically informed therapy; or 3) referral to a psychological or pain management specialist. Participants in the intervention group will additionally have access to an interactive website that provides information about whiplash and recovery relative to their risk category. For more information <https://www.mywhiplash.com.au/node/6>

Focus area 4 - (sub)acute low back pain

Imaging for low back pain

- Many imaging (radiography, CT scan, and MRI) findings identified in people with low back pain are also common in people without such pain, and therefore their importance in diagnosis is uncertain.(14, 30)
- **No evidence exists that imaging improves patient outcomes** and guidelines consistently recommend against the routine use of imaging for people with low back pain.(14, 30) Clinicians need to consider whether the **overall clinical picture** might indicate a serious cause for the pain onset, or the course, of low back pain ('red flags').(14)

Management

- International guidelines(30, 62-64) for patients with non-specific LBP recommend **simple first line care** (advice, reassurance and self-management) and a review at 1–2 weeks. If patients need second line care, non-pharmacological treatments (eg, physical and psychological therapies) are recommended before pharmacological therapies. If pharmacological therapies are used, the guidelines recommend the lowest effective dose and for the shortest period of time possible. Exercise and/or cognitive behavioural therapy, with multidisciplinary treatment for more complex presentations, are recommended for patients with chronic low back pain. Electrotherapy, traction, orthoses, bed rest, surgery, injections and denervation procedures are not recommended for patients with non-specific low back pain.
- The recent Lancet series describes the evidence to support interventions to improve the prevention and management of low back pain.(13)
 - The authors identified **a gap related to secondary prevention.**
 - The authors identified that there is evidence to support education and advice to remain active as the first-line of treatment for people with acute low back pain

- The authors identified evidence to support NSAIDs, superficial heat, massage, spinal manipulation or acupuncture as the second-line or adjuvant treatment for people with acute low back pain; and exercise therapy and cognitive behavioural therapy in selected patients.
- The authors acknowledge the lack of evidence for sub(acute) low back pain but suggest a reasonable approach is to use therapies for chronic back pain (multidisciplinary treatment)
- The **stepped approach** to the management of people with low back pain begins with more simple care that is progressed if the patient does not respond.
- **Other approaches** involve the early identification and management of psychosocial risk factors ('yellow flags') for progression to chronic, disabling pain. Two approaches include:
 - A stratified care approach whereby patients are stratified according to risk level (high, medium and low), and assigned treatment appropriate to the risk level.
 - A matched care approach, which is conceptually similar to the stratified approach, but aims to match the treatment to the specific risks.
- The **relative effectiveness of the stepped, stratified and matched care approaches** have not been evaluated.(31)
- A recent narrative review acknowledges that while the "wait and see" (stepped) approach is attractive given the limited time and resources available in primary care, it misses the opportunity for early intervention based on known risk factors of poor prognosis.(31) **For more information** about the advantages and disadvantages of the stepped, stratified and matched care approaches see this recent narrative review by Linton et al (2018).
- Another recent narrative review also highlights **the importance of identifying patients based on their specific characteristics** and suggests that combining patient's psychosocial profile with the activity-related behavioural style may be of added value in tailoring the patient's treatment to his/her specific needs.(65)
- A recent systemic review about the secondary prevention of chronic musculoskeletal pain (that included mainly low back pain patients) reported that compared with a "one-size-fits-all" treatment, **stratified programs** showed significant improvements in several domains of the International Classification of Functioning, Disability and Health.(36) *See 3.1 Options related to Goal 1- Focus area 1: multi-modal, not condition specific.*
- Multidisciplinary biopsychosocial rehabilitation for subacute low back pain has been reported in a recent Cochrane systematic review to be no better than a brief clinical intervention including education and advice about exercise **in interventions not targeted to high risk patients.**(35)
- **One example of a trial that implements risk-stratified care** for low back pain in family practices in the UK is the **IMPACT Back study**.(47, 66) The study reported significant improvements in patient disability outcomes and a halving in time off work, without increasing health care costs. The STarT Back tool is specifically designed for primary care settings and is used in the study as the subgrouping tool that allocates patients into low-, medium- or high-risk subgroups in order to guide decision making about treatment and referral. In the study, the tool is offered to GPs in both a computer-based format and a paper-based format completed by the GP and patient in the consultation. The targeted treatments include a minimal intervention delivered by GPs (for those patients at low risk of poor outcome) or referral to primary care physiotherapists who can apply physiotherapy approaches to addressing pain and disability (for those at medium risk) and additional cognitive-behavioural approaches to help address psychological and social obstacles to recovery (for those at high risk). *For information about the key content covered in the training packages for primary care providers see Appendix 10.*

- There is growing interest in the stratified care approach with two multi-site trials currently being conducted using a stratified care approach to prevent the progression to chronic pain in high-risk patients. *For more information see below: relevant research currently being conducted.*
- **One example of the matched care approach is the recent Australian trial (WISE study).**⁽¹⁹⁾ See 3.1 Options related to Goal 1- Focus area 5: work-related injuries.

For information about screening tools see 3.2.1 Face-to-face and/or online education and training.

ACI model of care for acute low back pain

- The **NSW Agency for Clinical Innovation (ACI) has recently developed a model of care (MOC) for acute low back pain (2016)** that can be applied to primary care and emergency department settings.⁽²⁸⁾ The ACI model of care was developed in collaboration with policy-makers, clinicians, consumers and researchers.
 - The model provides different care pathways according to a classification based on a diagnostic triage (acute or chronic non-specific low back pain, low back pain with leg pain and suspected serious spinal conditions).
 - Risk stratification is recommended (using tools such as the STarT Back or Orebro) to guide the amount and type of treatment provided
 - Follow-up reviews are recommended to monitor individuals' progress.
 - The key principles of the ACI model of care for acute low back pain include: *Principle 1- Assessment: history and examination (red flags); Principle 2 - Risk stratification (yellow flags); Principle 3 - Patient education; Principle 4 - Active physical therapy encouraged; Principle 5 - Begin with simple analgesic medicines; Principle 6 - Judicious use of complex medicines; Principle 7- Cognitive behavioural approach; Principle 8 - Only image those with suspected serious spinal pathology; Principle 9 - Predetermined times for review; and Principle 10 - Timely referral and access to specialist services.*
 - Psychologically informed practice⁽⁵⁶⁾ is recommended including teaching patients simple coping methods, helping patients to understand the relationship between beliefs and behaviours, and developing goal-orientated plans.
 - Referral of patients for more complex psychological intervention (e.g. clinical psychologist, physiotherapist trained in psychologically informed physiotherapy) is recommended for patients with 'yellow flags' or patients who develop chronic pain.
 - The MOC also includes language to avoid and language to use when speaking with patients about pain.
 - A consumer resource has been developed by the ACI based on this MOC.

Relevant research currently being conducted

- (US) Targeted interventions to prevent chronic low back pain in high-risk patients: A multi-site pragmatic cluster randomized controlled trial (**TARGET Trial**) <http://www.targettrial.pitt.edu/> This trial compares treatment for preventing transition to chronic low back pain via guideline-based primary care versus primary care plus timely referral to physical therapists trained in psychologically informed practice (PIP). Acute low back pain patients at all clinics are risk stratified (high, medium, low) using the STarT Back Tool.
- (Australia) Implementation of a clinical PATHway of CarE to improve patient health outcomes and reduce costs for common musculoskeletal disorders (low back pain, neck pain or whiplash, knee osteoarthritis) in primary care (**PACE study**). *For more information see above Focus area 1- multi-modal or not condition specific.*

Aboriginal and Torres Strait Islander people

- In-depth interviews with Aboriginal men and women with chronic low back pain in regional and remote areas of Western Australia **found the experience of chronic low back pain impacted on activities of daily life, employment, sport and family participation, emotional and cultural well-being.**(67)
- The interviews also identified that most participants **held biomedical beliefs** about the cause of low back pain, attributing pain to structural/anatomical vulnerability of their spine. This belief was attributed to the advice from healthcare practitioners and the results of **spinal radiological imaging**. The authors report that findings are consistent with research in other populations and infer that disabling chronic low back pain may be at least partly iatrogenic.(68)
- Health practitioners need to consider communication content and style to improve interactions with Aboriginal people with chronic low back pain.(69) Barriers to communication between health care providers and Aboriginal people with low back pain include communication content, information that was not evidence-based, miscommunications, communicative absence and the use of medical jargon. Enablers related to communication style described as **'yarning'**, a two-way dialogue, and healthcare practitioners with good listening and conversational skills.
- A qualitative randomised crossover design compared low back pain information (entitled **My Back on Track, My Future** [MBOT]) with an evidence-based standard (the Back Book [BB]).(70) Twenty Aboriginal adults from one rural region in Western Australia participated.
 - My Back on Track, My Future was developed as five short audio-visual scenarios, filmed using Aboriginal community actors. The project aimed to reduce the burden of low back pain amongst Aboriginal people in WA's Midwest and was a partnership between WACRH, the Geraldton Regional Aboriginal Medical Service (GRAMS), and Curtin University.
<http://www.wacrh.uwa.edu.au/my-back-on-track-my-future>

Key messages—My Back on Track, My Future

1. Back pain—it affects mind, body, and spirit (This message reflected the need to promote a biopsychosocial awareness of low back pain; and had a particular focus on managing anger and depressed mood, known to be a priority from our prior work)
 2. Permanent back damage is rare-big medical terms do not usually mean serious damage that can't improve (This refers to the misperceptions about low back pain perpetuated by misinterpreted radiological imaging, including terms such as "arthritis" and "degeneration")
 3. Moving well will help your back naturally (Referring to the need to encourage normal movement to improve low back health)
 4. Get your blood pumping (Referring to the benefits of exercise in the management of low back pain disorders in addition to general health and well-being)
 5. Look after yourself to look after your community (Referring to putting low back pain care as a priority; this responds to the tendency, particularly by some Aboriginal women, to prioritize the well-being of their family above their own health needs)
- **Results of the study:** thirteen participants preferred MBOT, four the BB, two both, and one neither. Participants valued seeing "Aboriginal faces," language that was understandable, the visual format, and seeing Aboriginal people undertaking positive changes in MBOT. In contrast, many participants found the language and format of the BB a barrier. Participants who preferred the BB were more comfortable with written information and appreciated the detailed content.

Focus area 5 – Return to work and work-related injuries

Management

- Models of care for injured workers in the compensation environment recommend the adoption of **biopsychosocial management approaches**. Compensation Models of Care following musculoskeletal

workplace injuries consistently recommend early diagnostic triage, identification of potential psychosocial obstacles to recovery, provision of education about pain biology, encouragement to keep active and encouragement to remain at work or an early return to work even when symptoms persist.(18)

- A recent systematic review highlighted that **multidisciplinary and multi-factorial interventions that seek to address a range of individual and societal factors** that influence return to work for workers with musculoskeletal and pain-related conditions are more likely to be effective.(71)
 - The review reported that graded activity programs need to be supplemented by workplace modifications (e.g. modified duties, modified working hours, supernumerary replacements, ergonomic adjustments) and/or service coordination (communication within the workplace or communication between healthcare providers and the workplace) to be effective in reducing lost time associated with work disability. These multi-component interventions also have a positive effect on improving work functioning after return to work and reduce costs associated with work disability.
- Psychologically-informed treatments provided for injured workers without psychological risk factors, are no better than usual treatment.(72)
- A recent Australian trial (WISE study) evaluated whether **an early intervention for high risk injured workers** was associated with fewer lost work days over two years compared to the usual (stepped) care.(19) Workers in the intervention were screened within 1–3 weeks of injury as being at high risk of delayed returned to work using the Örebro Musculoskeletal Pain Screening Questionnaire-short version (ÖMPSQ-SF) and were offered psychological assessment and a comprehensive protocol implemented by a range of stakeholders (including psychologists, physiotherapists, GPs, insurance case managers and return to work coordinators) working in collaboration, as needed, to address the identified psychological and workplace obstacles for return to work. The study found that at the two-year follow-up, the mean lost workdays for the intervention group was less than half of the usual care group, their claim costs were 30% lower, as was the growth trajectory of their costs after 11 months.

For information about screening tools see 3.2.1 Face-to-face and/or online education and training.

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain


- **Face-to-face multidisciplinary consumer pain programs for consumer with chronic pain** are currently implemented in six PHNs and WAPHA.
- In August 2019, **NT PHN** implemented a multidisciplinary pain management program for people with non-compensable chronic pain <https://www.painnt.com.au/>

3.1.3 Consumer initiative related to safe and effective use of medications and tapering of opioids

- A consumer initiative could be implemented by PHNs about simple analgesics, safe and effective use of medications and tapering of opioids
- A face-to-face consumer initiative about opioids could include:
 - Group based sessions (education and active, practice-based learning), perhaps embedded in a face-to-face multidisciplinary consumer pain program, or a separate consumer workshop/education session(s); or

- Individual sessions with a primary care provider such as a GP, practice nurse, community pharmacist to provide education and behavioural strategies (e.g. identifying practical and psychological barriers to tapering opioids and problem-solving solutions, non-pharmacological options, cognitive and behavioural strategies and goal setting); and
- Or an online consumer initiative about opioids (e.g. webinar, YouTube video) could include education and behavioural strategies.
- And Information resources (paper-based, links to online resources).
- Initiative could be tailored to specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Zang 2019- systematic review post-surgery (hospital setting)
- Mathieson 2019 systematic review -primary care setting (under review)- chronic pain
- Sullivan 2017- chronic pain
- Holliday 2017 –chronic pain
- Darnall 2019- perioperative digital intervention-breast cancer patients
- White 2016- YouTube

PHN initiatives

- Opioid Early Intervention Pilot Project (West Vic PHN)

Narrative synthesis

Focus area 6 - Opioid consumer initiatives

Consumer initiatives about opioids

- Research about consumer initiatives related to opioid use, non-initiation and tapering in people with acute and subacute pain in the post-surgery, post-injury and (sub)acute back pain setting, **particularly in the primary care setting is limited.**(25)
- A recent systematic review identified **strategies implemented in the hospital setting to reduce postsurgical opioid prescribing at discharge** including two patient interventions.(22)
 - One of the patient interventions examined the impact of preoperative patient education about opioids. The study found that only 10% of patients who had education sessions requested opioids after surgery compared with 100% in the control group and patients also experienced a lower intensity and duration of pain postoperatively.
 - The other patient intervention in this review examined the impact of a shared-decision making model. The study used a computer-based decision aid to help patients determine the amount of opioid they would be prescribed after caesarean delivery. The shared decision-making session resulted in a significant decrease in the amount of opioid prescribed. Postintervention, 8% of patients obtained a refill prescription and 90% of patients were satisfied or very satisfied with their pain management.

- A recent systematic review identified interventions **in the primary care setting related to opioid deprescribing involving patients with chronic pain**. The review found only a small number of trials related to opioid deprescribing that involved patients with chronic pain in the primary care setting.(73)
 - A consumer intervention identified in this systematic review is provided as **an example** of an initiative about tapering opioids:
 - The pilot RCT explored the effects of a tapering opioid intervention with patients recruited from specialist pain clinics and primary care clinics. The intervention included firstly, a Motivational Interviewing-based session with the patient concerning opioid tapering including: 1) eliciting the patient’s history related to pain, opioid therapy, and related difficulties; 2) eliciting change talk related to tapering; 3) education about dose-related health risks; 4) identifying practical and psychological barriers to tapering opioid dose and problem-solving ways to overcome these; and 5) developing a commitment to change with respect to opioid therapy. Patients were also shown a short video of interviews with the same patients who were in the first video concerning coping with challenges of tapering off opioids.
 - The taper support intervention protocol included an additional 17 weekly 30-minute individual sessions.
 - Patients were provided with an opioid medication prescription for the week at each visit.
 - Patients were encouraged to attend all sessions in person but were offered to complete up to every other session by telephone.
 - At the sessions, patients reported on pain, withdrawal, and mood/anxiety symptoms.
 - Each session included pain self-management training modelled after empirically supported cognitive-behavioural therapy (CBT) interventions for chronic pain. The sessions included: (1) rationale for pain self-management and education about the neurophysiology of pain and the role of cognitive and behavioural variables in chronic pain and adjustment to it; (2) behavioural goal setting; (3) education about, training in, and practice of various relaxation techniques (diaphragmatic breathing, progressive muscle relaxation, body scans, applied relaxation); (4) behavioural activation techniques, activity scheduling, and instruction in activity pacing; (5) education regarding the role of cognitions in negative affective responses to pain and instruction in positive pain coping self-statements and distraction techniques; (6) sleep hygiene education; and (7) education about and training in ways to maintain gains, reduce the risk of pain flare-ups, and cope with pain flare-ups if they do occur.
 - Motivational Interviewing was used periodically to address ambivalence about tapering as needed.
 - Patients completed “personal action plans” at each session for home activities to perform between sessions (e.g., practice of relaxation techniques, personal goal-related activities).
 - The group that had taper support improved significantly more than usual care in self-reported pain interference, pain self-efficacy, and prescription opioid problems at 22 weeks. The authors concluded that this taper support intervention is feasible and shows promise in reducing opioid dose while not increasing pain severity or interference.(74)
 - A recent pilot RCT study aimed to assess the feasibility of **digital perioperative behavioural pain medicine intervention in breast cancer surgery (“My Surgical Success”)** and evaluate its impact on pain catastrophizing, pain, and opioid cessation after surgery.(75)

- The intervention involved a 90-minute pain psychoeducational video, a downloadable Personalised Plan for Surgical Success and a downloaded relaxation audiofile.
- Video content included information and skills to regulate cognition, emotion, and physiologic hyperarousal related to pain, including relaxation, thought reframing, and behaviours that modulate attention and counteract helplessness about pain. During the video, learners were guided to self-tailor and apply the information by completing their Personalized Plan for Surgical Success.
- The study found that the intervention was acceptable and feasible and significantly accelerated opioid cessation after surgery. The authors concluded that perioperative digital behavioural pain medicine may be a low-cost, accessible adjunct that could promote opioid cessation after breast cancer surgery.
- The **Brainman video series** developed by the Hunter Integrated Pain Service, Hunter New England Local Health District and Hunter New England and South Coast PHN were released on YouTube as a low-cost public health intervention. Each video used an evidenced informed script appropriate for low literacy and a cartoonist to provide matching images. The videos included 1) Understanding pain in less than 5 minutes, and what to do about it! (2013 Jan 15). 2) Understanding Pain: Brainman stops his opioids (2014 Oct 3) 3) Understanding Pain: Brainman chooses (2014 Oct 3). (76) The videos were also widely circulated through varied professional and consumer networks <https://www.aci.health.nsw.gov.au/ie/projects/brainman>

Relevant PHN initiatives

PHN initiatives related to the secondary prevention of chronic pain

- The NT PHN is currently developing '**Prescribing and Supplying Opioids for Acute Pain Top End Health Service (TEHS) Guideline**'. The purpose is:
 - To promote safe, consistent and appropriate prescribing of opioids for acute pain inpatients and at discharge from hospital to optimise patient outcomes in relation to pain management.
 - Outline an approach to treating acute pain using a multimodal analgesic approach based on the recommendation of TEHS Acute Pain Service (APS).
- Western Victoria PHN has contracted La Trobe University to undertake the **Opioid Early Intervention Pilot Project** in three pharmacies. The project is funded through the Pharmacotherapy Area-Based Network for 18 months and involves providing pharmacies with additional funding to consult with a patient when they are initially prescribed an opioid. The intervention provides education on expectations for pain management and options other than opioids to manage their pain. Patients will be followed up at one, three and six months.

3.1.4 Transitions of care / pre-surgery consumer initiative

- A pre-surgery consumer initiative could be implemented by PHNs and could include:
 - Individual sessions with a primary care provider such as a GP or practice nurse and could include:
 - Pre-surgery risk assessment for developing chronic pain after surgery
 - Education about pain, and pharmacological and non-pharmacological pain management
 - Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management;

- Or online consumer initiative (e.g. webinar, YouTube video) including education and behavioural strategies;
- And information resources (paper-based, links to online resources).
- Initiative could be tailored to specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Chou 2016-Clinical guidelines-pain
- Shug 2015 -Acute pain management scientific evidence
- Nicholls 2018- psychological treatments-surgery
- Darnall 2019- breast cancer surgery- digital behavioural intervention
- Archer 2016 – cognitive behavioural therapy- lumbar spine surgery

Narrative synthesis

Focus area 2 - surgery

Management

- **Optimal management of postoperative pain begins in the preoperative period.**
 - The US clinical practice guidelines related to the management of postoperative pain, 2016, recommend that optimal management begin in the preoperative period with an assessment of the patient and development of a plan of care tailored to the individual and the surgical procedure involved.(26)
- **Pre-operative and peri-operative psychological-interventions have the potential to reduce chronic postsurgical pain (CPSP).**
 - *Acute Pain Management: Scientific Evidence (2015)* by the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine highlights that training in coping methods or behavioural instruction prior to surgery reduces pain, negative affect and analgesic use.(3)
 - A recent systematic review(38) found preliminary evidence to support perioperative cognitive behavioural therapy-based interventions to reduce post-surgery pain intensity and disability.
 - A digital perioperative behavioural pain medicine intervention in breast cancer surgery has been found to be acceptable and feasible and significantly accelerate opioid cessation after surgery.(75)
 - Another recent trial found that a cognitive-behavioural based physical therapy (CBPT) program delivered six weeks after lumbar spine surgery to patients with preoperative high fear of movement improved outcomes. Patient who had CBPT had significantly greater decreases in pain


and disability and increases in general health and physical performance compared to the education group.(77) The CBPT program was delivered face-to-face and by telephone and focused on self-management, problem solving, cognitive restructuring and relaxation training (see www.spine-surgery-recovery.com for more information).

- **Preoperative patient education is recommended**, although there is limited evidence to support the mode and content of the education.(3, 26)
 - *Acute Pain Management: Scientific Evidence (2015)* by the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine highlights that the benefit of preoperative patient education on pain outcomes is inconsistent, however, specific pain education in specific surgical settings may result in decreased pain, opioid use and less healthcare utilisation.(3) The working group suggests that structured preoperative education may be better than routine information.
 - The US clinical practice guidelines related to the management of postoperative pain, 2016, recommends preoperative education. However, the guidelines report that there is insufficient evidence to determine the comparative effectiveness of different type of patient education which can range from single episodes of face-to-face instruction or provision of written materials, videos, audiotapes, or Web-based educational information to more intensive, multicomponent preoperative interventions including individualised and supervised exercise, education, and telephone calls.(26)

3.1.5 Transitions of care / post-surgery consumer initiative

- A post-surgery consumer initiative could be implemented by PHNs and could include:
 - Individual sessions with a primary care provider e.g. GP, practice nurse, community pharmacist:
 - Post-surgery risk assessment for developing chronic pain
 - A clinical medication review including medication reconciliation with active patient counselling
 - Education about pain and pharmacological and non-pharmacological pain management
 - Behavioural strategies such as coping methods, though-reframing, relaxation, goal setting and self-management
 - Follow-up at, for example, 2 weeks, 6 weeks and 12 weeks:
 - Or online consumer initiative (e.g. webinar, YouTube video) including education and behavioural strategies;
 - And information resources (paper-based, links to online resources);
- Initiative could be tailored to specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Hesselink 2012 - systematic review- to improve patient handovers from hospital to primary care
- Ensing 2015 systematic review- community pharmacists
- Bethishou 2019 systematic review- community pharmacists

PHN initiatives

- The Pain Prescribing on Discharge Working Group including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health

Narrative synthesis

Transitions of care

- **Effective discharge planning, communication between hospital and primary care providers and follow-up of patients is crucial** for effective pharmacological and non-pharmacological pain management and tapering of opioids.
- A systematic review(78) highlighted the following interventions to **improve patient handovers from hospital to primary care** including
 - Medication reconciliation (Medication reconciliation is the process of creating the most accurate list possible of all medications a patient is taking — including drug name, dosage, frequency, and route — and comparing that list against the physician's admission, transfer, and/or discharge orders, with the goal of providing correct medications)
 - Discharge planning; shared involvement in follow-up by hospital and community care providers
 - Health system support:
 - Electronic tools to facilitate quick, clear, and structured summary generation
 - Use of electronic discharge notifications
 - Web-based access to discharge information for general practitioners

For more information about health system support related to discharge from hospital see Section 3.3.2 Transitions of care / health system support initiatives.

- **Community pharmacists** may play an important role in improving continuity of care during transition from hospital to the community in collaboration with hospital clinicians and primary care providers, although more research is needed.(79, 80)
 - A systematic review to identify the optimal role for pharmacists in care transitions during and after hospitalisation recommends the implementation of multifaceted programs that combine medication reconciliation with active patient counselling and a clinical medication review.(79) The authors also recommend that pharmacists collaborate with other health care professionals to acquire information about the clinical background of patients.

Relevant PHN initiatives

PHN initiatives related to the secondary prevention of chronic pain

- The **Pain Prescribing on Discharge Working Group** including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health. The PHN role will be to support the development of health literacy tools that are appropriate for the Indigenous population (patients will be provided with information on discharge); and to support integration of care between hospitals and primary care.

3.1.6 Telehealth assisted allied health services

- Telehealth (telemedicine) could be implemented by PHNs for consumers with (sub)acute pain at risk of developing chronic pain
- Telehealth could be used to provide:
 - Individual sessions with an allied health practitioner such as a physiotherapist or clinical psychologist e.g. Telehealth in Murrumbidgee Local Health District, connecting a senior physiotherapist in Griffith (base site) to the patient and an allied health assistant in Hay (recipient site); and/or
 - Group-based activities (e.g. education, exercise)
- Initiative could be tailored to specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Egmond 2018 systematic review – physiotherapy with telerehabilitation
- Gentry 2019- telehealth group-based programs (range of conditions)
- Mariano 2019- group based program- chronic pain

PHN initiatives

- Outreach services (telehealth and visiting health care providers) connecting people in pain with pain specialists and other health providers- chronic pain

Grey literature

- The NSW ACI. Improving physiotherapy access using telehealth. Murrumbidgee Local Health District, Report, 2018
- NSW ACI. Chronic Pain -Telehealth Pilot Project Evaluation Report 2016

Narrative synthesis

Focus area 1: multifocal or not condition specific

Telehealth

- **Access to care can be increased by telehealth.**(29, 32, 81, 82)

- An **example of a telehealth service related to physiotherapy** is the telehealth service in Murrumbidgee Local Health District which connects a senior physiotherapist in Griffith (base site) to the patient and an allied health assistant in Hay (recipient site). An evaluation (82) of the **physiotherapy telehealth intervention** in Murrumbidgee Local Health District reported: a) improved access to physiotherapy services in Hay from 60% to 80% of the time (over 12 months from March 2015– March 2016); b) reduced travel time for patients as well as out-of-pocket costs from an estimated \$445 per to \$10 per physiotherapy consultation; and c) strengthened capacity of the local care teams, particularly the competencies of the allied health assistants to support best practice care.
- An **example of telehealth services related to chronic pain** are the telehealth services at the Children’s Hospital Westmead and Orange Hospital at the chronic pain clinics, established as part of a pilot study (2016).(81) This project was a collaboration between ACI, LHDs and specialty networks, Healthdirect Australia (HDA), and the Ministry of Health. A variety of models were used by both sites during the pilot, for example 1:1 patient support with GP in attendance for initial assessment; 1:1 patient support with GP in attendance for follow up ; 1:1 multi-disciplinary assessment or treatment in the patient’s home with usual correspondence back to the GP after the consultation; follow up by individual disciplines e.g. telecounselling, physio in home with local physiotherapist in attendance; and upskilling local allied health and medical practitioners. Telehealth was found to be a feasible and effective model to deliver chronic pain services into primary care and patients’ homes.
- A recent systematic review(83) showed that **group-based telehealth** (video teleconference groups) for a range of conditions is feasible and produces treatment outcomes similar to in-person treatment, with high participant satisfaction despite technical challenges. Preliminary findings suggest that online group cognitive behavioural therapy may be as effective in improving coping among persons with chronic pain as in-person groups.(84) Additional research is needed to identify optimal methods of video teleconference group delivery to maximise clinical benefit and treatment outcomes.
- The NSW Agency for Clinical Innovation (ACI) has recently developed a **telehealth in practice guide** (2019) which describes the range of models for telehealth services; enablers and barriers to implementation; and a **readiness assessment checklist** to support clinicians and clinic administrators to consider the key features to successful implementation of telehealth.(85)
- **Further information and support to implement telehealth** includes:
 - Australian Government, Department of Health, MBS Online: Telehealth: Specialist video consultations under Medicare
 - NHMRC Centre for Research Excellence in Telehealth Policy Digest
 - A Practical Guide to Knowledge Translation in Telehealth (2016) NHMRC Centre for Research Excellence
 - NSW Agency for Clinical Innovation (ACI) Chronic Pain Telehealth Toolkit (2015)
 - NSW Agency for Clinical Innovation runs a Virtual forum of the Telehealth Capability Interest Group
 - NSW Agency for Clinical Innovation report *Improving physiotherapy access using telehealth. Murrumbidgee Local Health District, 2018*
 - NSW Agency for Clinical Innovation. *Chronic Pain -Telehealth Pilot Project Evaluation Report 2016*
 - The Allied Health Telehealth Capacity Building Scoping Project 2015 (Queensland Health)
 - The Allied Health Telehealth education package including an online, on demand training package (Queensland Health)
 - Evaluation Resource Guide: Allied Health Telehealth Capacity Building Project 2016 (Queensland Health)
 - The Allied Health Telehealth Network providing email group, intranet and scheduled videoconference presentations (Queensland Health)

For more information about the above see **Section 4**.

Focus area 2 - surgery

Telehealth

- A recent systematic review of **physiotherapy with telerehabilitation after surgery** found that physiotherapy with telerehabilitation is feasible and improves Quality of Life in surgery populations, although the overall effectiveness on functional outcomes could not be determined.(33)

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain

- Four regional PHNs currently implement **outreach services** (telehealth and visiting health care providers) connecting people in pain with pain specialists and other health providers.

3.1.7 Digitally delivered care: online consumer pain program and mobile apps

Online consumer pain program

- An online consumer pain program could be implemented by PHNs for consumers with (sub)acute pain at risk of developing chronic pain with varying levels of telephone and/or email support from a primary care provider (e.g. psychologist, physiotherapist training in psychologically informed practice may ring participants once a week or may be available to answer emails only)
- Based on the same principles as face-to-face programs and use online modules to teach pain management information and support patients to develop their self-management skills and encourage non-pharmacological approaches (e.g. keeping active, pacing, challenging unhelpful thoughts, simple coping strategies, relaxation and stress management, sleep, return to work); help consumers set goals and undertake graded exercise; and provide information about simple analgesics and tapering of opioids
- An online consumer pain program could also be tailored to specific (sub)acute pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]; or tailored to specific groups, for example Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Mobile app for the post-surgery or post-injury phase or (sub)acute back pain

- A mobile app could be developed by PHNs for consumers with (sub)acute pain at risk of developing chronic pain
- A mobile app could also be tailored to specific (sub)acute pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]; or tailored to specific groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.
- A mobile app could provide information and strategies similar to an online consumer pain program (see above)

Supporting evidence

Online consumer pain program



Peer review literature

- Dear 2018, 2015 - The Pain Course- chronic pain- available online through MindSpot
- Schultz 2018 - This Way Up- chronic pain –available online through St Vincent's Hospital Sydney

Mobile apps



Peer review literature

- Machado 2016-systematic review- low back pain
- Reynoldson 2014 review – pain self-management
- McKay 2018 systematic review - health behaviour change (range of conditions)

Narrative synthesis

Focus area 1: multifocal or not condition specific

Digitally delivered care

- Access to care can also be increased by **digitally delivered health care** e.g. online intervention, mobile apps.(32)
- **Internet-delivered pain management programs** are usually based on the same principles as face-to-face programs and use online modules to teach pain management information and support patients to develop their self-management skills. These programs can be offered in clinician guided formats, where patients are provided weekly support throughout the program via telephone or email, or in more self-guided formats with little or no clinician contact.(86)
 - For example, The Pain Course, a free internet-delivered pain management program managed by Macquarie University (NSW) for people with chronic pain, aims to provide information that helps participants understand chronic pain and their symptoms, and teach cognitive and behavioural self-management skills to help reduce pain-related disability, anxiety and depression.(32, 86, 87)
<https://ecentreclinic.org/?q=PainCourse> <https://mindspot.org.au/pain-course>

- Another example is an online multidisciplinary pain management program for people with chronic pain, This Way Up, managed by St Vincent's Hospital in Sydney (NSW) and requires a referral from a clinician (GP, pain physician, GP, nurse or allied health practitioner) <https://thiswayup.org.au/how-we-can-help/courses/chronic-pain/>.(88)
- However, there is limited research about internet-delivered pain management programs specifically for people with (sub)acute pain (post-surgery, post-injury, (sub)acute low back pain).
- **Mobile apps** have been used for pain self-management for people with chronic pain.(89-91)
 - However, a study of mobile apps for pain self-management reported variation in app quality and a lack of user and clinician engagement in development of these mobile apps.(89)
 - There is also limited research about mobile apps that relate to the (sub)acute pain phase (post-surgery and post-injury). See *Focus area 4: (sub)acute low back pain* for a description of a recent systematic review of mobile apps for the self-management of low back pain.

Focus area 4 - (sub)acute low back pain

Digitally delivered care

- In the recent Lancet series, the authors suggest that rural and remote regions rehabilitation advice and support given **online**, combined with self-management, might be an option where internet access is available.(11)
- A recent systematic review of **mobile apps** for the self-management of low back pain found that generally app developers are selecting interventions that are endorsed by guidelines, although their quality is low.(91) The authors reported that there are many apps available for the self-management of low back pain, but their effectiveness in improving patient outcomes has not been rigorously assessed. The authors suggest that app developers need to work closely with healthcare professionals, researchers, and patients to ensure app content is accurate and evidence-based and the app is user-friendly and useful.

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain

- Four PHNs currently implement **online consumer distribution platforms and information portals** (e.g. GoShare, Health Resource Directory, PatientInfo)

3.1.8 Peer support group/network

- Peer support group/network could be implemented by PHNs for consumers with (sub)acute pain at risk of developing chronic pain
- A peer support group/network could be delivered face-to-face, online or via social media
- A peer support group/network could also be tailored to specific (sub)acute pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]; or tailored to specific groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



PHN initiatives

- Face-to-face peer support network -Adelaide PHN- chronic pain

Narrative synthesis

Relevant PHN initiatives


PHN initiatives that could be adapted to the secondary prevention of chronic pain

- Adelaide PHN is currently implementing a face-to-face **peer support network** with visiting health professionals from a range of disciplines.

3.1.9 Community awareness campaign

- A community awareness campaign could be implemented by PHNs to improve the community's understanding of pain; how to prevent chronic pain; promote self-management and non-pharmacological approaches to (sub)acute pain; and promote safe and effective use of medicines.
- A community awareness campaign could be delivered via social media, television, radio, print media or community events
- A community awareness campaign could also be tailored to specific (sub)acute pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]; or tailored to specific groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- White 2016- Brainman video series- chronic pain focused

PHN initiatives

- Support for Pain Revolution
- Brainman video series (could be further adapted) HNESCPHN, HIPS, HNELHD

Narrative synthesis

Relevant PHN initiatives

PHN initiatives related to the secondary prevention of chronic pain

- Two PHNs support the **Pain Revolution** (<https://www.painrevolution.org>) which runs community awareness and health professional education events in rural and regional areas of Australia. The initiative aims to increase knowledge, skills and local support to prevent and manage chronic pain.

PHN initiatives that could be adapted to the secondary prevention of chronic pain


- The community awareness initiative, **Brainman video series**, developed by the Hunter Integrated Pain Service, Hunter New England Local Health District and Hunter New England and South Coast PHN <https://www.aci.health.nsw.gov.au/ie/projects/brainman> could include other videos relevant to the post-surgery, post-injury and (sub)acute back pain population.

3.1.10 Promotion of relevant consumer resources

- PHNs could promote relevant consumer resources via consumer and health professional networks (events and newsletters), HealthPathways and online consumer distribution platforms and information portals.
- *Examples:*
 - PAIN-ED website
 - Painaustralia website
 - NSW Agency for Clinical Innovation (ACI) - Best practice care for people with acute low back pain
 - painHEALTH website
 - Brainman brief educational videos
 - Keele group - the IMPaCT Back study - online training (UK)
 - NPS MedicineWise consumer information
 - Whiplash Injury Recovery - a Self Help Guide to aid the recovery of people with a whiplash injury - Motor Accident Insurance Commission

*For more information about the above consumer resources and for additional consumer resources see **Section 3**.*

3.2 Options related to Goal 2 (health professional capacity building)



3.2.1 Face-to-face and/or online education and training

- Face-to-face and/or online education and training could be implemented by PHNs for primary care providers **relevant to patients with (sub)acute pain and specific (sub)acute pain populations [e.g. post-surgery, post-injury, (sub)acute back pain]**
- Education and training could be delivered via different modes e.g. webinar, online course, online platform (Project ECHO)
- Education and training should involve a range of primary care providers including practice nurses, allied health practitioners and community pharmacists as well as GPs (or provide separate education and training for different disciplines)
- Face-to-face and/or online education and training should active, practice-based learning about the following issues (see *Principles for the secondary prevention of chronic pain*):
 - **Explaining pain:** How to explain pain to patients and language to avoid/use
 - **Imaging:** When to use imaging and how to explain imaging results to patients
 - **Risk factors for the progression of acute pain to chronic:**
 - How to assess patients for risk factors for poor recovery to inform type of treatment ('yellow flags')
 - How to use specific screening tools in the context of clinician's own clinical reasoning
 - **Psychologically informed practice:**
 - How to help patients set goals and promote self-management, pacing and other non-pharmacological approaches
 - How to apply simple coping methods and behavioural strategies
 - Referring to allied health practitioners (e.g. physiotherapists, psychologists)
 - Additional training for physiotherapists- psychologically informed physical therapy
 - **Safe and effective use of medicines/opioid education:**
 - Appropriate opioid prescribing
 - Non-initiation and deprescribing of opioids
 - Non-opioid pain analgesic medicines
 - **Follow-up of acute pain patients to monitor patient progress:** Scheduling times for follow-up (e.g. at 2, 6 and 12 weeks)

- **Integration of care between hospital setting and primary care; and between primary care providers**
 - Communicating with hospital services and other primary care providers
 - Timely referral to specialist services (e.g. at 12 weeks) if appropriate
- **Return to work:**
 - Approach to patients
 - Capacity certification
 - Understanding the compensable environment
- Education and training could also be provided for working with **relevant population groups** including Aboriginal and Torres Strait Islander peoples, people from culturally and linguistically diverse backgrounds, people from rural and remote areas, older Australians, people with dementia, children and young people and other relevant groups. For example,
 - For working with Aboriginal and Torres Strait Islander people, provide education and training for primary care providers about 'clinical yarning'; and provide cultural safety training for primary care providers and support staff.

For a summary of the enablers to implementing education and training initiatives identified in the consultation with PHNs in Phase 1 of the Chronic Pain Project see **Appendix 6**.

Supporting evidence



Peer review literature

- Sowden 2012 (IMPACT Back Pain) training programme
- Kelly 2018 (StressModex) – acute whiplash- stress inoculation training
- Brunner 2013 Systematic review (sub)acute low back pain
- Beales 2019-implementation of screening tool – physiotherapy clinics -return to work
- Papapagorus 2018 - Return to work- flowchart for certification – primary care
- Kelly 2017 – clinical prediction rule – acute whiplash
- Hall 2018- Physiotherapist-delivered cognitive-behavioural interventions- low back pain
- Keefe 2018 systematic review - psychologically informed practice for pain management- chronic pain
- Cowell 2019 – cognitive functional therapy-chronic pain
- Furlan 2019; Frank 2015; Katzman 2019, 2014; Shelley 2017; Flynn 2017 (Project ECHO- chronic pain)

PHN initiatives

- Most PHNs provide chronic pain management education events
- As part of the consumer pain program in South Eastern NSW PHN, ACI supported facilitators to access webinar skills training in pain management; putting cognitive behavioural therapy skills into practice (PMRI)
- Western Victoria PHN has implemented Project ECHO Opioid Management Clinic and is currently in the initial phase of implementing Project ECHO for chronic pain

Narrative synthesis

- A recent systematic review about the secondary prevention of chronic pain highlights the need for **adequate education of health professionals** to screen patients for risk factors and integrate a biopsychosocial perspective into their care.(36)
- The *Principles for the secondary prevention of chronic pain* highlighted in this rapid review include **education and training of primary care providers** to address issues such as risk factors and screening; self-management, goal-setting, pacing and non-pharmacological approaches; simple coping methods and behavioural strategies; opioid deprescribing; referring to allied health practitioners; follow-up; and return to work approaches, capacity certification and the compensable environment (See *Principles for the secondary prevention of chronic pain*).
- However, **education and training related to the secondary prevention of chronic pain** (as part of professional development or more formal undergraduate and postgraduate courses for GPs, physiotherapists, practice nurses and other primary care providers) is limited, although education and training related to the secondary prevention of chronic pain may be embedded in chronic pain management education and training.
- There is also **limited research** about the acceptability, feasibility or effectiveness of education and training **related to the secondary prevention of chronic pain**
- Furthermore, trials of interventions with people with acute and chronic pain often provide **insufficient details about the training** involved to implement the intervention (e.g. treatment manuals, patient materials and provider training outlines).(92)

Screening tools

Overall, there is a paucity of research related to the **implementation of screening tools** in primary care.(93)

Surgery

- **Screening tools** are usually surgery-type specific.(2) One generic tool assesses the effect of 14 biomedical and psychosocial items that were derived from a systematic review of the CPSP risk factor literature.(2, 46)

(Sub)acute whiplash

- A **clinical prediction rule (CPR)** has been validated which can identify patients from the early acute post injury stage who are at high risk of poor recovery at 12 months. The tool can also identify those who are likely to fully recover and a third medium risk group who could either recover or develop chronic pain and disability.(49) However, a qualitative study evaluating the perceptions of physiotherapists, chiropractors and osteopaths of adopting clinical prediction rule found that clinicians' understanding and use of the tool was mixed. The authors suggest that further education is needed so that clinicians understand the purpose of the tool, how to use the tool in the context of their own clinical reasoning and how to communicate the results to patients.(50)

(Sub)acute low back pain

- The **STarT Back tool** has been used in primary care in the UK to identify low, medium or high risk of persistent disability in patients with low back pain.(47, 48) The **IMPACT Back study** is one of the few trials that describes the training of primary care providers (GPs and physiotherapists) to implement the screening tool.(47, 66) *For information about the key content covered in the training packages see Appendix 10.* The Keele group also provides online training and useful resources to use the tool in routine care. <https://startback.hfac.keele.ac.uk/training/>

Return to work and work-related injuries

- The 10-item Swedish scale (the **Orebro Musculoskeletal Pain Screening Questionnaire - Short-Form - OMPSQ-SF**) has been used with injured workers in Australia with pain at any site, and training (online and face to face) is available for its use.(19, 20)
- A **validation study of Örebro Musculoskeletal Pain Screening Questionnaire-short version (ÖMPSQ-SF)** to predict time to return to pre-injury work duties following a work-related soft tissue injury (regardless of body location) provides strong support for the use of the ÖMPSQ-SF in an applied setting for identifying those injured workers likely to have delayed return to work when administered within 15 days of the injury.(20)
- A recent pre-post study in Western Australia described the **implementation of the 10-item Örebro Musculoskeletal Pain Screening Questionnaire (ÖMPSQ-10) in private physiotherapy clinics in patients with a compensable musculoskeletal problem**.(93)
 - The authors described the key barriers to the use of screening questionnaires (from their experience) as: lack of time as perceived by clinicians; lack of clinician knowledge on the utility of screening questionnaires; and lack of clinician knowledge on how to use screening questionnaires to inform clinical decision making.
 - The study implemented clinician-focused strategies (group education, additional individual education if needed and supplementary written material) and organisation-focused strategies (paperwork/front office procedure for questionnaire completion and familiarisation of new employees). Education of physiotherapists emphasised that the questionnaire should be used to assist, rather than being the sole indicator used, in developing the individual client's management pathway.
 - The study reported a significant positive shift in behaviour to more frequent use of the OMPSQ-10 for new compensable patients.
- A recent Australian trial (WISE study) evaluated **an early intervention for high risk injured workers**.(19) Workers in the intervention were screened within 1–3 weeks of injury as being at high risk of delayed returned to work using the Örebro. *See 2.1 Options related to Goal 1 - Focus area 5: work-related injuries.*

Return to work and work-related injuries

- A recent systematic review found that graded activity programs need to be supplemented by service coordination (communication within the workplace or **communication between healthcare providers and the workplace**) and/or workplace modifications to effectively reduce lost time associated with work disability for workers with musculoskeletal and pain-related conditions.(71) *See 2.1 Options related to Goal 1- Focus area 5: work-related injuries.*
- 'Helpful' perspectives **for primary care providers to support recovery and return to work** are highlighted in a recent narrative review of the management of musculoskeletal pain in a compensable environment(18) and include understanding that work is critical for good health and a therapeutic intervention rather than solely an outcome and that many workers return to work before they have recovered 100%. The authors also highlight the need for **education for healthcare practitioners** about compensation-based system issues.
- A recent review provides a systematic approach for Australian primary care providers (including general practitioners and physiotherapists) **to support return to work through appropriate certification**.(21)
 - The paper provides a return to work flowchart to support systematised and appropriate certification of capacity developed by the Transport Accident Commission and WorkSafe Victoria.
 - The authors suggest the following practice point: *Certification should relate to the injured worker's capacity to safely undertake tasks at work as it relates to the injury, rather than other*

workplace issues. Notably, certifying 'unfit' may not necessarily resolve the incorrect assumption or the situation limiting return to work. For example, if a clinician deems that the injured worker is 'unfit' and has no capacity to work, then consider what they would be doing at home. For example, if they are bed-bound, then it is likely that certifying 'unfit' is medically necessary, whereas outside this circumstance then there are various levels of capacity that could facilitate return to work. Asking 'What would the person reasonably be doing at home?' is a key question in determining someone's functional capacity and whether certifying 'unfit' is medically necessary.

- The authors suggest a more systematised approach to certification coupled with professional education and support may reduce variations and inaccuracies in certification, improve return to work rates and reduce the increasing burden of disease related to workplace injuries.
- The negative impact of long-term work absence, work disability and unemployment on the individual and their family following injury is acknowledged in a recent editorial.(94) The editorial suggests that **physiotherapists promote a positive relationship between work, rehabilitation and health** to patients to reduce progression to “worklessness” following injury by following strategies such as:
 - a) Ask your patient about their work before they were injured or unable to work and document the tasks they were doing; b) Start a conversation early about work, including setting a date with the injured person about when you expect s/he will be able to be back at work; c) Create an expectation that work is part of rehabilitation, not the end result; d) Dispel the myth that a person needs to be back to normal or pain free to work; e) Communicate clearly with all stakeholders about your role, scope and limitations; you can't find someone a job, but you can remove some barriers; f) Acknowledge and promote your expertise in physical health and functioning, liaise with the workplace and the medical practitioner to facilitate return to pre-injury or modified work duties; g) If you work in a jurisdiction where physiotherapists have the legal authority to certify physical capacity, consider doing so; h) Reflect on your words and avoid catastrophic language so that you do not create fear and avoidance in your patients around movement, activity and work; and i) Avoid becoming embroiled in system bashing or clinician criticism, which creates further anger, distrust and perceptions of injustice in your patients.

Psychologically informed practice

- There is a growing interest in **psychologically informed practice for pain management** and primary care physicians and physical therapists have delivered psychologically informed practice as part of a stratified care approach involving screening and targeting of treatment for people at high risk for pain-associated disability.(56)
- In a recent narrative review of psychologically informed practice, the authors describe **the major types of psychological interventions** as: a) educational (threat reduction and activation); b) behavioural change (explicit focus on incorporating adaptive behaviours in response to pain); c) cognitive-behavioural (principal focus on cognition and coping strategies); d) psychophysiological focus (variants of stress reduction and mindfulness); and e) contextual cognitive-behavioural therapy (acceptance and commitment therapy).(56)
- A recent systematic review reported that **physiotherapist-led cognitive-behavioural interventions** (that included techniques designed to reduce fear of movement and pain-related disability such as pacing, goal setting, problem-solving, relaxation, and challenging unhelpful thoughts relevant to low back pain) were effective for low back pain.(92)
 - The review highlighted that using a cognitive-behavioural approach, including a variety of techniques that could be easily adopted in a physical therapy setting, provides greater benefits for patient outcomes compared to brief education, exercise or physical techniques (such as manual therapy) alone.

- The review found that with additional training, physiotherapists can deliver effective cognitive-behavioural interventions.
- However, the authors report that without training or resources, successful translation and implementation remains unlikely. The authors suggest that researchers improve reporting of procedural information, provide relevant materials, and offer accessible provider training.
- However, a recent systematic review found that **physiotherapists lack confidence** in their ability to identify, communicate and manage cognitive, psychological and social dimensions of chronic low back pain in practice.(95)
 - Physiotherapists report feeling that neither their initial training nor currently available professional development equipped them to successfully deal with these factors in practice and **emphasised a need for training** on integrating these factors into patient management.(95, 96)
 - The authors of a recent narrative review of psychologically informed practice suggest the following 3-step training approach: a treatment manual, an experiential workshop, and ongoing supervision with consultation and feedback.(56)
- **Cognitive functional therapy (CFT)** is a behaviourally oriented intervention that targets patients' individual biopsychosocial profiles.(96, 97)
 - A recent qualitative study of a 10-month formal CFT training programme, which included a combination of didactic learning, problem-based learning, communication training, video reviews and direct feedback on clinical practice, has the capacity to enhance physiotherapists' self-reported confidence and competence in managing the biopsychosocial dimensions of nonspecific chronic low back pain.(96) The authors suggest that to achieve an enduring change in clinical behaviour, biopsychosocial training should include clinical integration and ongoing support.
- **Stress inoculation training and exercise** has been recently implemented in a trial for patients with acute whiplash-associated disorder at risk of poor recovery.(10)
 - In the study, a physiotherapist provided six sessions (one per week) to patients of stress inoculation training, teaching strategies to assist participants in managing acute stress responses. It consisted of three phases: (1) identify and understanding stress-identifying specific stressors and how these affect pain, behaviour, emotions, physical performance and thoughts; (2) developing skills for managing stress, such as relaxation, problem solving and helpful coping self-statements; and (3) applying skills in various stressful situations to develop tolerance and confidence. Participants were encouraged to practise these skills on a weekly basis with home practice.
 - The training for physiotherapists to conduct stress inoculation training with patients included a two-day training workshop conducted by a clinical psychologist, rehabilitation physician and musculoskeletal physiotherapist.
 - Physiotherapists reported that the training developed their confidence to deliver the program and supported using at least components of the program in routine practice.(98) The authors conclude that physiotherapists with some additional training, could apply stress inoculation training for whiplash-injured patients at risk of poor recovery.

Research currently being conducted

- A US multi-site pragmatic cluster randomized controlled trial (**TARGET Trial**) is currently being conducted to compare treatment for preventing transition to chronic low back pain via guideline-based primary care versus primary care plus timely referral to physical therapists trained in psychologically

informed practice. *For more information see 2.1 Options related to Goal 1 - Focus area 4: (sub)acute low back pain.*

Project ECHO

- **Project Extension for Community Health Outcomes (ECHO)** was originally developed by the University of New Mexico's Health Science Centre to build the capacities of primary care providers and to increase access to specialist care in rural and underserved populations.(99)
- Project ECHO expands primary care provider capacity to manage complex diseases by sharing knowledge, disseminating best practices, and building a community of practice. The model has expanded rapidly with over 140 ECHO projects currently (in 2018) established globally.(100)
- Project ECHO uses a **"Hub" and "Spoke" model** to promote knowledge exchange between health care specialists typically located at academic centres ("the hub") and primary care providers (PCPs) at the front line of community health care (the "spokes"). Through regularly scheduled virtual educational clinics, Project ECHO creates a supportive community network where PCPs can connect with specialists and with other PCPs practicing in similar settings via multipoint video technology to discuss best practices in care and complex cases managed within their practice. This multifaceted knowledge and capacity-building intervention includes two fundamental components: a didactic lecture delivered by a member of the hub team (based on curriculum developed from guidelines, best practices, and/or a needs assessment); and recommendations for case management (telementoring) offered by the community in response to anonymised clinical cases presented by spoke sites.(100)
- Project ECHO has been implemented **to address chronic pain**.(101-106)
- A systematic review of the impact of Project ECHO on participants and patient outcomes has shown it to be an effective and potentially cost-saving model.(107)
- Key factors found to increase registrant engagement include relevant and practical curriculum content; strong and supportive relationships among learners, ECHO faculty, and workplace colleagues; and innovative learning approaches that included opportunities for active, virtual participation through technology, participant management activities, and ECHO's unique curriculum design.(108)

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain


- Most PHNs currently implement **education and training for primary care providers** related to chronic pain management.
- **Western Victoria PHN has implemented Project ECHO Opioid Management Clinic.** The Project ECHO Opioid Management Clinic is currently available on a weekly basis as an online one-hour meeting connecting addiction medicine specialists and psychiatrists at St Vincent's hospital (hub) with primary care providers (spoke) to upskill primary care providers in opioid management. The program is funded through the Pharmacotherapy Area Based Networks.
- Western Victoria PHN is currently in the initial phase of implementing **Project ECHO for chronic pain**.

3.2.2 Opioid initiative about prescribing, non-initiation and deprescribing of opioids

- Implementation of an initiative for primary care providers about opioids including prescribing, non-initiation and deprescribing of opioids including face-to-face or online (e.g. webinar or Project ECHO)

- An opioid initiative could include:
 - Education and active, practice-based learning e.g. workshop, short course, online module; Project ECHO; and/or
 - Behavioural strategies e.g. clinical reminder system, audit and feedback
- Initiative could be tailored to working with specific population groups, for example, Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Zhang 2019 systematic review Behavioural Interventions to Decrease Opioid Prescribing After Surgery
- Stanley 2019 - Australian hospital opioid prescribing initiative
- Holliday -GPs trainees-chronic pain
- Mathieson 2019 systematic review -primary care setting- chronic pain (under review)

PHN initiatives

- Prescribed Drugs of Dependence Active Learning Modules - Western Victoria PHN
- Project ECHO (opioid management) – Western Victoria PHN
- Some PHNs provide education and training events about opioids or embedded in chronic pain management events

Narrative synthesis

Opioid initiatives for clinicians

- **Research about health professional initiatives related to opioid use, non-initiation and tapering** in people with acute and subacute pain in the post-surgery, post-injury and (sub)acute back pain setting, particularly in the primary care setting, **is limited**.
- A recent systematic review identified **strategies to reduce postsurgical opioid prescribing at discharge**, implemented in the hospital setting.(22) Six types of behavioural interventions were identified: local consensus-based processes (18 studies), patient-mediated interventions (2 studies), clinical practice guidelines (1 study), educational meetings (1 study), interprofessional education (1 study), and clinician reminder (1 study). All but one study reported a statistically significant decrease in the amount of opioid prescribed at discharge after surgery, and only 2 studies reported evidence of increased pain intensity.
- A recent systematic review identified interventions **in the primary care setting related to opioid deprescribing involving patients with chronic pain**.(73) The review found only two trials that focused on interventions for clinicians, one trial evaluated training sessions and decision tools; and the other trial evaluated online education of patient simulation plus case-based learning.
- **An example** of opioid deprescribing intervention in an Australian hospital:
 - A recent Australian study implemented in an orthopaedic specialty unit in St Vincent's Hospital, Melbourne examined the impact of an intervention comprising (i) an Expert Advisory Group

oversight of opioid prescribing, (ii) development of a prescription opioid guideline for various hospital contexts and (iii) a series of education sessions.

- The pre-post study found a significant reduction in the number of patients discharged on combination opioids from 71.4% to 45.7%, a significant reduction in the provision of full pharmaceutical quantities of opioid on discharge from 29.4% to 6.1% and a significant increase in opioid weaning plans included in discharge summaries from 6.9% to 87.4%.(109)
- **An example** of an education intervention about opioids for Australian general practitioner trainees:
 - A recent Australian study investigated the impact of an educational intervention for general practitioner trainees emphasising limitations, risk mitigation, and deprescribing of opioids with transition to active self-care. This educational intervention incorporated pre-readings, a resource kit, and a 90-minute interventional video case-based workshop incorporated into an education day.
 - The pre-post study found that the education intervention produced significant changes to trainees' judgments about, and intentions toward, long-term opioid analgesia maintained at two months.(110)
 - *For more information about the content of the education initiative see **Appendix 11**.*

Relevant PHN initiatives

PHN initiatives related to the secondary prevention of chronic pain

- One PHN has implemented an education and training initiative related to opioid use. The **Prescribed Drugs of Dependence Active Learning Modules** aims to improve risk management and treatment pathways for patients being prescribed drugs of dependence in general practice. The program supports practices to embed a consistent approach to quality and safer prescribing of drugs of dependence. The program consists of three face-to-face workshop sessions for GPs, practice nurses, pharmacists and allied health and practice support to improve opioid prescribing and pain management in primary care. The sessions are accredited for the Royal Australian College of General Practitioners CPD points. The role of the PHN is to coordinate speakers and running of these events. *Enablers:* practice visits to promote the program resulted in program enrolments; and incentives funded by the Pharmacotherapy area-based network (Ballarat region – Western Victoria PHN) whereby GPs receive \$1000 if they can demonstrate (using a template to collect information) that they have improved their opioid prescribing.
- **Western Victoria PHN has implemented Project ECHO Opioid Management Clinic (see above).**

3.2.3 Interdisciplinary community of practice

- Implementation of a face-to-face and/or online interdisciplinary community of practice (CoP) to improve clinical practice, build relationships, promote knowledge-sharing and referral pathways
- CoP could include:
 - Chronic pain CoP or as part of a mental health COP or Alcohol and Other Drugs (AOD) CoP; and/or
 - A network for primary care providers involved in a specific face-to-face consumer pain program

Supporting evidence



PHN initiatives

- One PHN (Gold Coast) and WAPHA provide a network for primary care providers involved in a face-to-face consumer pain program

Narrative synthesis

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain

- Two PHNs currently provide a **network for primary care providers involved in a face-to-face consumer pain program for people with chronic pain.**

3.2.4 Promotion of relevant education and training and resources offered by other agencies

- Promotion of relevant education and training and resources for primary care providers offered by other agencies (e.g. face-to-face workshop or short course, online modules, postgraduate training)
- Promoted via health professional networks (events and newsletters) and HealthPathways
- *Examples include:*
 - NSW Agency for Clinical Innovation (ACI) Management of people with acute low back pain (2016)
 - NPS Medicine Wise e.g. Taking Action for Acute Low Back Pain - Online Case Study, Chronic Pain - Opioids and Beyond- Clinical case study
 - Better Pain Management online program - Faculty of Pain Medicine (FPM) and the Australian and New Zealand College of Anaesthetists (ANZCA)
 - Pain Management Research Institute, University of Sydney: Webinar skills training in pain management: putting cognitive behavioural therapy skills into practice
 - Annual Multidisciplinary Pain Management Workshop (1 week) by the Pain Management Research Institute, University of Sydney (PAINRefresh)
 - Keele group - IMPaCT Back study (STarT Back tool)- online training (UK)
 - The Australian Physiotherapy Association Level 1 course specifically targeting screening in the acute pain phase and how to prevent chronic pain
 - Pain Revolution
 - Curtin University School of Physiotherapy and Exercise Science
 - The University of Sydney Master of Medicine (Pain Management)
 - Royal Australian and New Zealand College of Radiologists (RANZCR) Educational modules (EMs) have been developed to improve the appropriateness of referrals for medical imaging

- Royal Australian College of General Practitioners (RACGP):
 - Webinar to equip GPs with the knowledge to help patients re-establish their health through active self-management and managing common complicating aspects of chronic pain
 - Active learning modules: Cognitive behavioural therapy skills for general practice (not pain specific)
 - Active learning modules: Psychological strategies skills training (not pain specific)
 - Assessment and Management of Acute Pain - Interactive Online Module

*For more information about the above and for additional education and training opportunities and resources offered by other agencies see **Section 4**.*


Narrative synthesis

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain

- As part of the Chronic Pain Management Program (consumer pain program) in South Eastern NSW PHN, the NSW Agency for Clinical Innovation (ACI) supported consumer pain program facilitators to access webinar skills training in pain management. Pain Management Research Institute, University of Sydney: **Webinar Skills Training in Pain Management: Putting Cognitive Behavioural Therapy Skills into Practice:** Online, interactive webinars training in CGT skills for pain: seven sessions over 2-3 months <https://sydney.edu.au/medicine-health/our-research/research-centres/pain-management-research-institute/postgraduate-and-short-courses-in-pain-management/putting-cognitive-behavioural-therapy-skills-into-practice.html> This webinar skills training would be relevant to the secondary prevention of chronic pain.
- **PHNs promote education and training and other resources for primary care providers offered by other agencies about chronic pain** via health professional networks (events and newsletters). Some of this education and training and other resources would also be relevant for the secondary prevention of chronic pain. *For more information about the above and for additional education and training opportunities and resources offered by other agencies see **Section 5**.*

3.3 Options related to Goal 3 (health systems support initiatives)



3.3.1 Implementation of relevant referral pathways

- Implementation of HealthPathways by PHNs to assist general practitioners (GPs) with the management of patients and the referral of patients to specialists and allied health professionals.
- Develop **relevant referral pathways to assess and manage patients with (sub)acute pain (including post-surgery, post-injury and (sub)acute low back pain populations)** including referral pathways to allied health practitioners and specialist services (if required at 12 weeks)
- Provide information about risk factors and screening tools; self-management, goal-setting, pacing and non-pharmacological approaches; simple coping methods and behavioural strategies; prescribing, deprescribing and non-initiation of opioids; follow-up; and return to work (see *Principle 4: Education and training of primary care providers*)
- **Provide clinicians with links to relevant education and training** and other health professional resources
- **Enablers** highlighted in recent research include: establishing workgroups (with GPs, specialists, allied health professionals) to create a sense of community and momentum, a forum for identifying system and service level issues and key insights and as a way of disseminating information; involving a range of clinicians in implementation to enable 'buy-in' including senior clinicians and executive staff; focusing on GPs that are new to the district (including visits and training); utilising existing PHN training events; and thinking about how to engage clinicians that are outside the normal engagement channels.
- Researchers also highlight that patients and clinicians do not think in terms of PHN boundaries and there is a need to create access to different pathways outside PHN boundaries.
- *For a summary of the enablers to implementing Healthpathways identified in the consultation with PHNs in Phase 1 of the Chronic Pain Project see **Appendix 8**.*

Supporting evidence



Peer review literature

- Gill 2019- South-West Victoria HealthPathways (not pain specific)
- Stokes 2018 - New Zealand HealthPathways (not pain specific)
- Gray 2018 Hunter and New England HealthPathways (not pain specific)

PHN initiatives

- HealthPathways - Most PHNs have implemented HealthPathways

Grey literature

- Norris 2018 Sydney HealthPathways (not pain specific)

Narrative synthesis

Co-ordination and continuity of care

HealthPathways

- HealthPathways is a password-protected website that contains information designed to assist general practitioners (GPs) with the management of patients and the referral of patients to specialists and allied health professionals (111, 112). The overarching objective of HealthPathways is to ensure that the right patient is referred to the right place at the right time and with the right information.
- A recent evaluation of the **Hunter and New England HealthPathways** identified that the critical elements for acceptability, growth and sustainability are the strong relationships between primary care and specialist clinicians, as well as formal partnerships that are built from the processes used to develop HealthPathways.(111) The authors recommend an approach to pathway development that engages GPs and specialists using a team-based process to build relationships and gain acceptance and endorsement of the pathways. Implementation that is accompanied by service redesign was found to be more likely to result in improvements. However, the authors acknowledge that this is a challenge due to the resourcing required.
- Recent evaluations of **HealthPathways in New Zealand**(113) and **South-West Victoria**(114) reported that clinicians viewed HealthPathways positively and usage had increased over time but there was a lack of planning and engagement of clinicians and that many GPs "did not think to look at HealthPathways or simply did not know about it."(114)
- **HealthPathways Sydney** was recently evaluated by the Menzies Centre for Health Policy, University of Sydney.(112) The evaluation recommended strategies to improve the planning of referral pathways and engagement of clinicians such as:
 - Establishing workgroups (with GPs, specialists, allied health professionals) to create a sense of community and momentum, a forum for identifying system and service level issues and key insights and as a way of disseminating information
 - Involving a range of clinicians in implementation to enable 'buy-in' including senior clinicians and executive staff

- Focusing on GPs that are new to the district (including visits and training)
- Utilising existing PHN training events; and thinking about how to engage clinicians that are outside the normal engagement channels.
- The authors also highlight that patients and clinicians do not think in terms of PHN boundaries and there is a need to create access to different pathways outside PHN boundaries.

Relevant PHN initiatives

PHN initiatives that could be adapted to the secondary prevention of chronic pain

- **Implementation of HealthPathways** to assist general practitioners (GPs) with the management of patients and the referral of patients to specialists and allied health professionals.
 - Most PHNs have implemented HealthPathways
 - Some PHNs have developed relevant referral pathways to assess and manage patients with (sub)acute pain (including post-surgery, post-injury and (sub)acute low back pain populations)

3.3.2 Transitions of care / health system support initiative

- A health system support initiative related to discharge from hospital could be implemented by PHNs to support better co-ordination of care between hospital and primary care
- Health systems support initiatives could include, for example:
 - Electronic tools to facilitate quick, clear, and structured summary generation, use of electronic discharge notifications; and online access to discharge information for general practitioners.
 - Discharge planning with shared involvement in follow-up by hospital and community care providers
- Initiative could be tailored to specific population groups, for example Aboriginal and Torres Strait Islander people, people from culturally and linguistically diverse backgrounds.

Supporting evidence



Peer review literature

- Hesselink 2012 - systematic review- to improve patient handovers from hospital to primary care
- Ensing 2015 systematic review- community pharmacists
- Bethishou 2019 systematic review- community pharmacists

PHN initiatives

- The Pain Prescribing on Discharge Working Group including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health

Narrative synthesis

- For information about pre-surgery consumer initiatives see Section 3.1.4 Transitions of care / pre-surgery consumer initiative; and for information about post-surgery consumer initiatives see Section 3.1.5 Transitions of care / post-surgery consumer initiative.

Transitions of care

- **Effective discharge planning, communication between hospital and primary care providers and follow-up of patients is crucial** for effective pharmacological and non-pharmacological pain management and tapering of opioids.
- Research on methods and outcomes of discharge planning and improving patient handovers from hospital to primary care are limited.(26, 78)
- The US clinical practice guidelines related to the management of postoperative pain, 2016, recommend a coordinated approach to discharge instruction including advice from hospital-based prescribers, nurses, physiotherapists, and pharmacists.(26)
- A systematic review(78) highlighted the following interventions to **improve patient handovers from hospital to primary care** including
 - Medication reconciliation
 - Electronic tools to facilitate quick, clear, and structured summary generation
 - Discharge planning; shared involvement in follow-up by hospital and community care providers
 - Use of electronic discharge notifications
 - Web-based access to discharge information for general practitioners

Relevant PHN initiatives


PHN initiatives related to the secondary prevention of chronic pain

- The **Pain Prescribing on Discharge Working Group** including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health. The PHN role will be to support the development of health literacy tools that are appropriate for the Indigenous population (patients will be provided with information on discharge); and to support integration of care between hospitals and primary care.

3.3.3 Electronic Persistent Pain Outcomes Collaboration (ePPOC)

- Electronic Persistent Pain Outcomes Collaboration (ePPOC), a national benchmarking system for the pain sector, could support the evaluation of a consumer pain program for consumers with (sub)acute pain at risk of developing chronic pain.

Supporting evidence



Peer review literature

- Tardiff 2016

PHN initiatives

- ePPOC reports for individual PHNs and pain services

Narrative synthesis

Benchmarking system for the pain sector

- The **Electronic Persistent Pain Outcomes Collaboration (ePPOC)** is a national benchmarking system for the pain sector (<https://ahsri.uow.edu.au/eppoc/index.html>) (115) It aims to improve clinical outcomes for people experiencing persistent pain through reporting and benchmarking. It is implemented and managed by the Australian Health Services Research Institute (AHSRI), University of Wollongong. It includes ePPOC for adults and PaedePPOC for children.
- **Multidisciplinary pain services** (public and private) involved in the ePPOC initiative routinely collect data using validated assessment tools; submit data to ePPOC every 6 months; and receive feedback and biannual reports. Services can compare their outcomes with the Australasian average and ePPOC benchmarks.
- Multidisciplinary pain services are currently mainly hospital based but there is a growing involvement of primary care services with several PHNs currently involved in the ePPOC initiative.

Relevant PHN initiatives


PHN initiatives that could be adapted to the secondary prevention of chronic pain

- Some PHNs are currently participating in the **Electronic Persistent Pain Outcomes Collaboration (ePPOC)** (national benchmarking system for the pain sector <https://ahsri.uow.edu.au/eppoc/index.html>) to evaluate a face-to-face multidisciplinary consumer pain program for consumers with chronic pain.

3.3.4 Prescription drug monitoring systems

- PHNs could provide education and training to support for the implementation of prescription drug monitoring systems e.g. SafeScript

Supporting evidence



PHN initiatives

- The Victoria PHNs led by Western Victoria PHN have been commissioned to provide education and training (in partnership with NPS MedicineWise) for GPs and pharmacists to support the implementation of SafeScript.

Grey literature

- Drugs and Poisons Information System Online Remote Access (DORA)
- Chronic Pain MedsCheck Trial

Narrative synthesis

Drug monitoring

- **SafeScript** is a real-time prescription monitoring and clinical decision support system that aims to provide doctors and pharmacists access to an up-to-the-minute medication supply history for certain high-risk medicines for their patient at the point of consultation. This includes all Schedule 8 medicines and other high-risk medicines such as benzodiazepines, zolpidem or zopiclone, quetiapine and codeine. It aims to help prescribers and pharmacists to safely manage patients who may be misusing prescription medicines, or those who may be receiving supplies of high-risk medicines beyond therapeutic need.
 - The Victorian Government has engaged Western Victoria PHN as lead for a consortium comprising all Victorian PHNs and NPS MedicineWise, to develop and deliver training for doctors and pharmacists to ensure successful uptake of the system. This includes safe and appropriate prescribing of high-risk medicines; drug counselling skills and engaging in conversations with patients around prescription medicine misuse and tapering of prescription medicines; and how information in the SafeScript system may be used to inform clinical decisions and regulatory obligations. Western Victoria PHN have undertaken a pilot of the training (face-to-face sessions and online NPS resources), which is now being rolled out <https://vtphna.org.au/safescript/> <https://vtphna.org.au/safescript-training-hub/>
- **Drugs and Poisons Information System Online Remote Access (DORA)** is a real-time prescription monitoring and clinical decision support system implemented by the Department of Health in Tasmania. <https://www.dhhs.tas.gov.au/psbtas/publications/general/dora>
- The **Chronic Pain MedsCheck Trial** is funded by the Australian Department of Health as part of the Sixth Community Pharmacy Agreement (6CPA) Pharmacy Trial Program (PTP). The 6CPA PTP was established to trial new and expanded Community Pharmacy Programs that seek to improve clinical outcomes for consumers and by progressing the role of community pharmacies in the delivery of primary healthcare services. The trial intervention from community pharmacy includes:
 - Supported self-management of patients taking medication who are dealing with chronic pain for more than three months through pharmacist advice
 - Pharmacy-based evaluation of patient's medicine
 - Provision of an action plan - action plans will incorporate education, self-management and referral to other health professionals where additional support is required
 - Three month follow up after the initial service

The final patient recruitment and initial service deadline has now been extended until 30 November 2019 and final patient follow up services will be completed 3 months later by the end of February 2020.
<http://6cpa.com.au/pharmacy-trial-program/chronic-pain-medscheck-trial/>

Relevant PHN initiatives

- (As stated above) Support for the implementation of **prescription drug monitoring systems, e.g. SafeScript**. The Victoria PHNs (N=6) led by Western Victoria PHN have been commissioned to provide education and training (in partnership with NPS MedicineWise) for GPs and pharmacists to support the implementation of SafeScript. Western Victoria PHN has undertaken a pilot of the education and training (face-to-face sessions and online NPS resources) which is currently being rolled out across Victoria.

3.3.5 Other quality improvement systems

Relevant PHN initiatives

- The **Practice Incentives Program (PIP)**, introduced in August 2019 by the Department of Health, Australian Government is a payment to general practices that participate in quality improvement activities to improve patient outcomes, and deliver best-practice care. There are 8 individual incentives under 3 payment streams. Most payments are for quality care activities, including for eHealth, quality improvement, teaching, Indigenous health, after hours care, procedural activities and rural locations.
- Information, guidelines, online forms and online training to help manage participation in the PIP is provided by the Department of Human Services, Australian Government.
<https://www.humanservices.gov.au/organisations/health-professionals/services/medicare/practice-incentives-program>
- The PIP Eligible Data Set is collected against 10 specified improvement measures from the Clinical Information Systems (CIS) of participating practices and **is submitted to the local PHN quarterly**. The Australian Institute of Health and Welfare (AIHW) is the National Data Custodian for the PIP Eligible Data Set.
- The PIP could potentially drive quality improvement activities e.g. health professional education and training, transitional care initiative *related to the management of patients with acute and subacute pain (e.g. post-surgery, post-injury and (sub)acute low back pain)*.

4. Consumer resources relevant to the secondary prevention of chronic pain

Examples of online consumer information

- **NSW AGENCY FOR CLINICAL INNOVATION(ACI), BEST PRACTICE CARE FOR PEOPLE WITH ACUTE LOW BACK PAIN**, a consumer resource to support the *Management of people with acute low back pain (2016)* model of care https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0003/363450/ALBP-Consumer-Info.pdf
- **NSW AGENCY FOR CLINICAL INNOVATION (ACI) PAIN MANAGEMENT NETWORK**, website designed to help consumers gain a better understanding of their pain. The site contains information to enable consumers to develop skills and knowledge in the self-management of pain in partnership with their healthcare providers. Key pages include 'Pain management for everyone' and 'PainBytes for youth'. <https://www.aci.health.nsw.gov.au/chronic-pain/for-everyone> <https://www.aci.health.nsw.gov.au/chronic-pain/painbytes>
*Note, website focuses on chronic pain but information is also relevant to (sub)acute pain
- **PAINHEALTH** aims to help health consumers with musculoskeletal pain access reliable, evidence-based information and tips to assist in the co-management of musculoskeletal pain. The website provides several online information modules, including 'Movement with pain' and 'Work related pain'. painHEALTH is an initiative of the Department of Health, Western Australia. <http://painhealth.csse.uwa.edu.au/>
Slater H, Davies S, Milne G et al. The painhealth website: A Western Australian policy-into-practice initiative to deliver holistic, consumer-focused best-evidence pain management for people with musculoskeletal pain. Physiotherapy (United Kingdom). 2015;1:eS1410.
- **PAIN-ED** is an online resource for patients and healthcare practitioners regarding evidence-based management of pain. It was developed by clinical researchers who recognised the need to translate the scientific evidence about pain for both public and healthcare practitioners. The site aims to dispel some common myths about chronic pain and provide hope for change. PAIN-ED is available at: <http://www.pain-ed.com/>
- **PAINAUSTRALIA** is Australia's leading pain advocacy body working to improve the quality of life of people living with pain, their families and carers, and to minimise the social and economic burden of pain on individuals and the community. The website provides information for consumers and health professionals. <https://www.painaustralia.org.au/>
- **BRAINMAN BRIEF EDUCATIONAL VIDEOS:** 1) Understanding pain in less than 5 minutes, and what to do about it! 2013 Jan 15. www.youtube.com/watch?v=C_3phB93rvI 2) Understanding Pain: Brainman stops his opioids. 2014 Oct 3 www.youtube.com/watch?v=Ml1myFQPdCE 3) Understanding Pain: Brainman chooses. 2014 Oct 3 www.youtube.com/watch?v=jlwn9rC3rOI
- The **KEELE GROUP - THE IMPACT BACK STUDY** evaluating risk-stratified care for low back pain in family practices in the UK provides patient information leaflets <https://startback.hfac.keele.ac.uk/patients/>
- **NPS MEDICINEWISE CONSUMER INFORMATION** NPS MedicineWise has a range of consumer education pages related to pain management, including '10 things you need to know about low back pain'. Freely available online. <https://www.nps.org.au/consumers/10-things-you-need-to-know-about-low-back-pain>
- **WHIPLASH INJURY RECOVERY – A SELF HELP GUIDE** to aid the recovery of people with a whiplash injury- Motor Accident Insurance Commission. NHMRC Centre of Research Excellence in Recovery Following Road Traffic Injuries <https://cre-rftrt.centre.uq.edu.au/whiplash-evidence-based-resource/whiplash-injury-recovery-%E2%80%93-self-help-guide>

- **AUSTRALIA ACCEPTANCE & COMMITMENT THERAPY WORKSHOPS & TRAINING** is a form of cognitive-behavioural treatment called acceptance and commitment therapy (ACT) used in the Toronto General Hospital Transitional Pain Service (TPS) ACT Groups & Centres Australia & New Zealand <https://www.actmindfully.com.au/>
- **MYBACKPAIN WEBSITE AND RESOURCES** provide information on back pain. The resource was developed through a collaboration between The University of Queensland, Arthritis Australia and the Cochrane Back and Neck Group. <https://mybackpain.org.au/>
- **THE ACUTE PAIN MANAGEMENT SERVICE, METRO NORTH HOSPITAL AND HEALTH SERVICE – PATIENT INFORMATION SHEET** 'Patient Controlled Analgesia (PCA) Breaking the Pain Barrier - Your pain relief in hospital', QLD Health. <https://metronorth.health.qld.gov.au/rbwh/wp-content/uploads/sites/2/2017/06/patient-controlled-analgesia.pdf>
- **MUSCULOSKELETAL AUSTRALIA (MSK) ONLINE RESOURCES AND COMMUNITY WEBINARS**, provides information to help consumers manage their condition. Recording from webinars are available online free, including one Webinar recording 'Understanding and managing low back pain' (21 August 2019) also available on YouTube. <https://www.youtube.com/watch?v=Cv9vteCdVMo&feature=youtu.be>
<https://www.msk.org.au/back-pain/>
<https://www.msk.org.au/community-webinars/>
- **THE AMERICAN CHRONIC PAIN ASSOCIATION (ACPA) COMMUNICATION TOOLS**, developed to help patients talk more productively with their health care provider or pharmacist about their pain. The tools can also help to identify patterns in patients' daily life that may have an impact on pain. Available freely online for consumers and healthcare providers. <https://www.theacpa.org/conditions-treatments/conditions-a-z/acute-pain/>
*Note, website focuses on chronic pain but information is also relevant to (sub)acute pain
- **PERSISTENT PAIN HUB, BETTER HEALTH CHANNEL INFORMATION ON PAIN AND PAIN MANAGEMENT**, including links to information and resources for people with pain. Provided by the Victorian Department of Health. Information has been translated into multiple languages. Freely available online for people experiencing pain. <https://www.betterhealth.vic.gov.au/conditionsandtreatments/pain>
*Note, website focuses on chronic pain but information is also relevant to (sub)acute pain
- **MY BACK ON TRACK, MY FUTURE**
LBP information (entitled My Back on Track, My Future [MBOT]) was developed as five short audio-visual scenarios, filmed using Aboriginal community actors. This project aimed to reduce the burden of LBP amongst Aboriginal people in WA's Midwest and was a partnership between WACRH, the Geraldton Regional Aboriginal Medical Service (GRAMS), and Curtin University. The information below was developed to help Aboriginal people manage their low back pain. <http://www.wacrh.uwa.edu.au/my-back-on-track-my-future>

Examples of support (Face-to-face, online, and/or telephone)

- **MUSCULOSKELETAL AUSTRALIA (MSK) ONLINE RESOURCES AND HELPLINE**. Helpline nurses and volunteers take calls from patients who need help managing back pain and other musculoskeletal conditions. Resources are also available online for various musculoskeletal conditions. National helpline phone 1800 263 265 or email helpline@msk.org.au. <https://www.msk.org.au/back-pain/>
*Note, uncertain to what extent this initiative offers support for consumers with (sub)acute pain
- **AUSTRALIAN PAIN MANAGEMENT ASSOCIATION (APMA) SUPPORT GROUPS**, aim to offer positive support and encouragement, with a focus on supporting and encouraging self-management of pain and use of evidence-based treatment, APMA currently operates a national network of Pain Support Groups (PSG) throughout Australia, which meet regularly. The groups are for everyone in the community living with pain (and their family members) and are FREE for APMA members. <https://www.painmanagement.org.au/what-we-do/support/pain-support-groups.html> *Note, uncertain to what extent this initiative offers support for consumers with (sub)acute pain

5. Health professional education and training and other resources relevant to the secondary prevention of chronic pain

Examples of information and support to implement telehealth

- **Australian Government, Department of Health, MBS Online: Telehealth: Specialist video consultations under Medicare** <http://www.mbsonline.gov.au/internet/mbsonline/publishing.nsf/Content/mbsonline-telehealth-landing.htm>
- **The NHMRC Centre for Research Excellence in Telehealth Policy Digest** <https://cretelehealth.centre.uq.edu.au/policy-digest> includes information about existing telehealth policies, position statements, guidelines, frameworks and standards. The aim of the Telehealth Policy Digest is to support health care practitioners and health service delivery organisations when they are considering, setting up or extending telehealth services and wish to develop their own policies, processes and standards.
- **A Practical Guide to Knowledge Translation in Telehealth (2016)** www.cretelehealth.org.au
- **The NSW Agency for Clinical Innovation (ACI) Chronic Pain Telehealth Toolkit (2015)** [https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0006/297843/Chronic Pain Telehealth Toolkit.pdf](https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0006/297843/Chronic_Pain_Telehealth_Toolkit.pdf)
- **The NSW Agency for Clinical Innovation (ACI)** runs a **Virtual forum of the Telehealth Capability Interest Group** on the third Thursday of the month and provides the opportunity for two presentations that highlight the use of telehealth in clinical practice. <https://www.aci.health.nsw.gov.au/make-it-happen/telehealth> Information on the Telehealth Capability Interest Group www.aci.health.nsw.gov.au/make-it-happen/telehealth/telehealth-capability-interest-group
- **NSW Agency for Clinical Innovation (ACI). Guidelines for the use of Telehealth for Clinical and Non Clinical Settings in NSW** Chatswood: ACI; 2015. Available from: https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0010/258706/ACI-telehealth-guidelines.pdf Guidelines for the use of Telehealth for Clinical and Non Clinical Settings in NSW www.aci.health.nsw.gov.au/_data/assets/pdf_file/0010/258706/ACI-telehealth-guidelines.pdf
- **NSW Agency for Clinical Innovation (ACI) Improving physiotherapy access using telehealth.** Murrumbidgee Local Health District 2018 - A telehealth extension to the existing physiotherapy service was introduced, connecting a senior physiotherapist in Griffith (base site) to the patient and an allied health assistant in Hay (recipient site). The report describes the benefits, key elements, services and physiotherapy patient flow in Hay; building engagement with primary healthcare; governance, planning and resourcing; and monitoring and evaluation https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0019/406018/Improving-physiotherapy-telehealth.pdf
- **The NSW Agency for Clinical Innovation (ACI). Telehealth in practice Guide (2019)** https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0008/509480/ACI_0261_Telehealth_guidelines.pdf
- **NSW Agency for Clinical Innovation (ACI). Telehealth contacts in NSW** https://www.aci.health.nsw.gov.au/_data/assets/pdf_file/0006/509478/Telehealth-contacts-in-NSW.pdf
- **The Allied Health Telehealth Capacity Building Scoping Project** is a joint initiative of the Allied Health Professions' Office of Queensland (AHPOQ), Health Service and Clinical Innovation Division, Department of Health and the Cunningham Centre, Darling Downs Hospital and Health Service (DDHHS).

<https://www.health.qld.gov.au/cunninghamcentre/html/telehealth> Nielsen I, Kirkpatrick J. Allied Health Telehealth Capacity Building: Scoping Project Completion report. Brisbane: Queensland Health; 2015. Available from: https://www.health.qld.gov.au/_data/assets/pdf_file/0020/150149/telehealthreportpt1.pdf

- **The Allied Health Telehealth education package.** The Cunningham Centre has developed an online, on demand training package available to all Queensland Health staff. This package is designed to increase clinician capability in the use of telehealth for the delivery of allied health clinical services in Queensland Hospital and Health Services. Further details regarding this package are listed below.
<https://www.health.qld.gov.au/cunninghamcentre/html/telehealth#objective1>
- **The Allied Health Telehealth Network** commenced in 2015 and provides members with opportunities to share information and develop collaborative partnerships through three main methods:
 - Opt-in email group coordinated by the Cunningham Centre for dissemination of information on telehealth collaboration, education / training, information sharing opportunities
 - Intranet/internet page for publishing of key documents such as information on networking structures, presentation schedule, summary information from previous presentations (AHPOQ or Cunningham Centre) [in development]
 - Scheduled videoconference presentations on allied health telehealth services implemented in Queensland HHSs (and other agencies if relevant).
<https://www.health.qld.gov.au/cunninghamcentre/html/telehealth#objective2>
- **Evaluation Resource Guide: Allied Health Telehealth Capacity Building Project** Published by the State of Queensland (Queensland Health), September 2016.
<https://www.health.qld.gov.au/cunninghamcentre/html/telehealth#objective3>
An electronic version of this document is available at
<https://www.health.qld.gov.au/ahwac/html/publications.asp>

Examples of education and training and other health professional resources for primary care providers

- **PAIN MANAGEMENT RESEARCH INSTITUTE (PMRI), UNIVERSITY OF SYDNEY: WEBINAR SKILLS TRAINING IN PAIN MANAGEMENT: PUTTING COGNITIVE BEHAVIOURAL THERAPY SKILLS INTO PRACTICE.** Online, interactive webinars training in CGT skills for pain: seven sessions over 2-3 months <https://sydney.edu.au/medicine-health/our-research/research-centres/pain-management-research-institute/postgraduate-and-short-courses-in-pain-management/putting-cognitive-behavioural-therapy-skills-into-practice.html>

As part of the Chronic Pain Management Program (consumer pain program) in South Eastern NSW PHN, the NSW Agency for Clinical Innovation (ACI) supported consumer pain program facilitators to access webinar skills training in pain management.
- **FACULTY OF PAIN MEDICINE, AUSTRALIAN AND NEW ZEALAND COLLEGE OF ANAESTHETISTS (ANZCA). BETTER PAIN MANAGEMENT: PAIN EDUCATION FOR PROFESSIONALS.** An online education program designed for specialist and general medical practitioners, medical students, nurses and allied health practitioners engaged in the care of patients with persistent pain. The program comprises twelve online education modules each designed to be completed in one hour. Relevant modules for (sub)acute pain available e.g. Module 1: Making an effective pain diagnosis: a whole person approach Module 2: The impact and management of psychological factors in pain Module 5: Identification and management of low back pain in the primary care setting Module 6: Opioids in pain management Module 9: Post-discharge acute pain management. Modules can be purchased individually for \$38.50 each or \$330 for the whole program.
<https://www.betterpainmanagement.com>
- **ANNUAL MULTIDISCIPLINARY PAIN MANAGEMENT WORKSHOP** (1 week) **PAIN MANAGEMENT RESEARCH INSTITUTE, UNIVERSITY OF SYDNEY.** PAINRefresh: Assessing and managing patients with acute and chronic pain conditions including sessions on screening and treatment options if identified early. Based on Sydney Medical School's internationally recognised postgraduate program in pain management, this workshop is

aimed at all health professionals interested in pain management. Topic areas include; concepts and assessment of pain, early intervention, opioid tapering, acute pain services etc. The workshop is being held on multiple dates between Mon 3rd February - Fri 7th February 2020, at Northside Conference Centre, Crows Nest. Registration costs \$930/\$1090 (early bird before November 1st /normal) and includes catering. <https://sydney.edu.au/medicine-health/our-research/research-centres/pain-management-research-institute/postgraduate-and-short-courses-in-pain-management/pain-management-multidisciplinary-workshop.html>

- **THE UNIVERSITY OF SYDNEY MASTER OF MEDICINE** (Pain Management) <https://sydney.edu.au/courses/uos-landing.html/content/courses/courses/pc/master-of-medicine-pain-management.html> Individual modules can be completed e.g. Acute Pain
- **PAIN MANAGEMENT RESEARCH INSTITUTE (PMRI) SYMPOSIUM, 'DESCENDING THE ANALGESIC LADDER: THE HOW, WHEN AND WHY OF OPIOID TAPERING FOR CHRONIC PAIN'**. A one-day symposium bringing together 13 national and international experts on the role of opioids in chronic pain management. Learning objectives include 1) Assess patients' appropriateness for opioid tapering and pain self-management; 2) Communicate the risks of opioid use for chronic pain and the benefits of tapering; 3) Explain pain self-management to patients. Held on 29 November 2019 9.00 – 18.00, Taronga Zoo, Sydney, Australia. Registration is AUD\$375 and includes catering for the day. Email paineducation.admin@sydney.edu.au for further information.
- **NSW AGENCY FOR CLINICAL INNOVATION (ACI) MANAGEMENT OF PEOPLE WITH ACUTE LOW BACK PAIN (2016)** which highlights three important areas for improvement: more appropriate clinical examination and use of radiological imaging only as necessary; better use of appropriate analgesia; and enhanced patient education. The model of care is designed for people presenting to health practitioners in primary care settings but is also a guide for care in settings such as emergency departments. <https://www.aci.health.nsw.gov.au/resources/musculoskeletal/management-of-people-with-acute-low-back-pain/albp-model>
- **NSW AGENCY FOR CLINICAL INNOVATION (ACI) PAIN MANAGEMENT NETWORK. ONLINE WEBINAR BURN PAIN PRESENTED BY SUE TAGGART.** 'Acute and chronic pain in the minor burn'. Available online free <https://vimeo.com/165508797>
- **NPS MEDICINE WISE CLINICAL RESOURCES ABOUT ACUTE NON-SPECIFIC LOW BACK PAIN** including research, factsheets, podcasts and professional development <https://www.nps.org.au/professionals/low-back-pain>
Taking Action for Acute Low Back Pain - Online Case Study - 2018 – 2019: This online interactive case study provides a clinical update on quality use of imaging and managing patients with low back pain. The activity emphasises the importance of using a risk stratification approach to identify risk factors for poor prognosis and inform selection of targeted and tailored interventions. It also includes resources to facilitate patient conversations on the importance of staying active during recovery.
- **VIC HEALTH AND NPS MEDICINEWISE** Management of acute non-specific low back pain. <https://www2.health.vic.gov.au/~media/Health/Files/Collections/Policies%20and%20guidelines/safe-opioid-use/Management%20of%20acute%20non-specific%20low%20back%20pain%20-%20for%20health%20professionals>
- **VIC HEALTH AND NPS MEDICINEWISE** Recommendations for deprescribing or tapering opioids <https://www2.health.vic.gov.au/~media/Health/Files/Collections/Policies%20and%20guidelines/safe-opioid-use/Recommendations%20for%20deprescribing%20or%20tapering%20opioids%20-%20for%20health%20professionals>
- **NPS MEDICINEWISE EDUCATIONAL VISITS** are facilitated by NPS MedicineWise Clinical Service Specialists to health professionals free of charge on a range of therapeutic areas commonly managed in general practice, including low back pain. <https://www.nps.org.au/cpd/book-a-visit>

- **NPS MEDICINEWISE NATIONAL PRESCRIBING CURRICULUM (NPC)** The National Prescribing Curriculum is a series of interactive case-based online modules that encourage confident and rational prescribing. Targeted at undergraduate and postgraduate medical students, Pharmacy, dental, nurse practitioner and other health professional students. One module on 'Analgesia for low back pain 2019' focuses on management of acute low back pain. The modules are free; however, users need a free NPS MedicineWise account to access them modules. <https://learn.nps.org.au/course/index.php?categoryid=78>
- The **KEELE GROUP - IMPACT BACK STUDY** evaluating risk-stratified care for low back pain in family practices in the UK also provide online training and useful resources <https://startback.hfac.keele.ac.uk/training/>
- The **AUSTRALIAN PHYSIOTHERAPY ASSOCIATION** Level 1 course specifically targeting screening in the acute pain phase and how to prevent chronic pain. <https://australian.physio/pd>
- **PAIN REVOLUTION:** Local Pain Educator to upskill rural health professionals with the latest pain science research and provide a support network to assist health professionals to share this knowledge with their local professional and public communities. <https://www.painrevolution.org/>
- **CURTIN UNIVERSITY SCHOOL OF PHYSIOTHERAPY AND EXERCISE SCIENCE-** undergraduate and postgraduate pain study units <https://healthsciences.curtin.edu.au/schools/physiotherapy-exercise-science/>
- **COURSES RUN BY PETER O'SULLIVAN, PROFESSOR OF MUSCULOSKELETAL PHYSIOTHERAPY AT CURTIN UNIVERSITY, PERTH, AUSTRALIA.** For example, Challenging common beliefs about pain. <http://www.pain-ed.com/>
- The **ROYAL AUSTRALIAN AND NEW ZEALAND COLLEGE OF RADIOLOGISTS (RANZCR)** Educational modules (EMs) have been developed to improve the appropriateness of referrals for medical imaging. <https://www.ranzcr.com/our-work/quality-standards/education-modules>
- The **ROYAL AUSTRALIAN COLLEGE OF GENERAL PRACTITIONERS (RACGP):**
 - <https://www.racgp.org.au/education/professional-development/online-learning/webinars/chronic-pain/managing-chronic-pain-in-general-practice> This webinar equips GPs with the knowledge to help patients re-establish their health through active self-management and managing common complicating aspects of chronic pain such as mental health (e.g. depression, anxiety, family problems). It discussed the benefits and harms of prescribing opioids and provides strategies for de-prescribing opioids. It is presented by Dr Simon Holliday and Dr Chris Hayes and developed in partnership with NSW Health.
 - RACGP active learning modules: Cognitive behavioural therapy skills for general practice (not pain specific) The ALM provides GPs with foundational CBT skills to use in their general practice and is ideal for GPs who have completed mental health skills training (MHST) and wish to include CBT in their clinical practice. <https://www.racgp.org.au/education/professional-development/online-learning/active-learning-modules/alm-cognitive-behavioural-therapy-skills>
 - Psychological strategies skills training (not pain specific). The RACGP Rural training package provides GPs with essential training in Focussed Psychological Strategies Skills Training (FPS ST) to enable rural GPs to provide CBT-derived FPS counselling to patients. <https://www.racgp.org.au/education/professional-development/online-learning/online-learning-modules/online-focussed-psychological-strategies-skills>
 - *Assessment and Management of Acute Pain - Interactive Online Module (8 hours). Provided by Learn EM,* this interactive online module reviews the approach to assessment and management of a patients' acute and ongoing pain management in various settings relevant to General Practitioners. Registration for this course requires a LearnEM account, and costs \$9.90. <https://moodle.learnem.com.au/enrol/index.php?id=498>
<https://www.racgp.org.au/education/courses/activitylist/activity/?id=58748&q=keywords%3dacute%2bpain%26triennium%3d17-19>
 - Approach to low back pain – exercise physiology. non-pharmacological treatments for back pain <https://www.racgp.org.au/download/Documents/AFP/2014/November/201411Clinical-Booth.pdf>

- **PAIN-ED** an online resource for patients and healthcare practitioners related to evidence-based management of pain. It was developed by clinical researchers who recognised the need to translate the scientific evidence about pain for both public and healthcare practitioners. The site aims to dispel some common myths about chronic pain and provide hope for change. PAIN-ED is available at: <http://www.pain-ed.com/>
- **PAINAUSTRALIA** is Australia's leading pain advocacy body working to improve the quality of life of people living with pain, their families and carers, and to minimise the social and economic burden of pain on individuals and the community. The website provides information for consumers and health professionals. <https://www.painaustralia.org.au/>
- **AUSTRALIA ACCEPTANCE & COMMITMENT THERAPY WORKSHOPS & TRAINING** is a form of cognitive-behavioural treatment called acceptance and commitment therapy (ACT) used in The Toronto General Hospital Transitional Pain Service (TPS) ACT Groups & Centres Australia & New Zealand <https://www.actmindfully.com.au/>
- **BOSTON CHILDREN'S HOSPITAL AND HARVARD MEDICAL SCHOOL PAEDIATRIC PAIN AND OPIOID EDUCATION' ONLINE ACCREDITED COURSE.** The online course provides training in acute and chronic pain management for paediatric patients, with an emphasis on safe and effective opioid use. Module 1 covers Acute Pain treatment; Module 2 covers Chronic Pain Treatment; and Module 3 covers Oversight and Safe Storage of Opioid Medications. Target audience includes Physicians and Trainees, Paediatric Physicians and Trainees, Paediatric Intensive Care (ICU) MDs and Trainees. The course takes 5 hours and 30 minutes to complete, and costs US\$125. Registration requires a free OPENPediatrics account. <https://www.openpediatrics.org/course/pediatric-pain-and-opioid-education-0>
- **THE HEALTH EDUCATION AND TRAINING INSTITUTE (HETI) NSW (2015). eLEARNING MODULE ACUTE PAIN MANAGEMENT FOR ADULTS.** This module uses a case study to demonstrate effective and appropriate pain management, including differentiating types of pain and how to relieve/manage acute pain. Target audience includes clinical nurses, midwives and junior medical officers. Registration requires a NSW Health account. <https://www.heti.nsw.gov.au/education-and-training/courses-and-programs/acute-pain-management-for-adults>
*Note, uncertain whether this course is still available
- **MUSCULOSKELETAL AUSTRALIA (MSK). HEALTH PROFESSIONAL WEBINAR 'DIAGNOSIS AND MANAGEMENT OF LOW BACK PAIN IN PRIMARY CARE'.** Presented by Dr Adrian Traeger, University of Sydney. This online webinar is designed to provide participants with clinically meaningful skills and knowledge to deliver recommended care. Recordings of individual webinars can be purchased online for \$30/\$45 (MSK member/non-member). <https://www.msk.org.au/health-professional-webinars>
- **WORKCOVER QUEENSLAND (2017). WEBINAR EARLY INTERVENTIONS FOR MUSCULOSKELETAL DISORDERS.** This Workers' Compensation Regulator webinar is presented by Michael Donovan who discusses; the background of musculoskeletal disorders, injury, pain and early intervention, the levels of prevention and intervention, and key components of successful intervention. Video recording available online free. <https://www.worksafe.qld.gov.au/forms-and-resources/webinars/early-interventions-for-musculoskeletal-disorders>
- **BRITISH PAIN SOCIETY AND THE FACULTY OF PAIN MEDICINE. E-PAIN: A MULTI-DISCIPLINARY PROGRAMME DESIGNED TO IMPROVE THE EARLY DIAGNOSIS AND MANAGEMENT OF PAIN.** The programme includes 12 training modules that cover knowledge ranging from how to manage acute pain well, through to learning in depth about common pain conditions and moves on to cover how to manage pain in specialist areas, like pain in cancer or pain in childhood. The modules are made up of 30-minute interactive e-learning sessions and assessments. For example, one session on Pharmacology for Acute Pain, explains the mechanisms of acute pain and its management using multimodal analgesia. Access to the e-PAIN programme is available from Australia and requires a free e-LfH account. Option to purchase the whole program (consisting of >60 sessions) for £90 or individual modules for £20. <https://www.e-lfh.org.uk/programmes/pain-management/>

- **OPENPEDIATRICS (2015) ONLINE VIDEO 'TREATMENT OF ACUTE PAIN IN A HEALTHY TEENAGER'**. The video demonstrates the clinical administration of pain treatment and opioid prescribing using the CRAFFT screening tool as a guide.
<https://www.openpediatrics.org/assets/video/treatment-acute-pain-healthy-teenager>

References

1. Lin I, Green C, Bessarab D. Improving musculoskeletal pain care for Australia's first peoples: better communication as a first step. *Journal of physiotherapy*. 2019;65(4):183.
2. Glare P, Aubrey KR, Myles PS. Transition from acute to chronic pain after surgery. *The Lancet*. 2019;393(10180):1537-46.
3. Schug SA, Palmer GM, Scott DA, Halliwell R, Trinca J. Acute pain management: scientific evidence. *Acute Pain Management: Scientific Evidence*. 2015:ixiv.
4. Katz J, Weinrib AZ, Clarke H. Chronic postsurgical pain: from risk factor identification to multidisciplinary management at the toronto general hospital transitional pain service. *Canadian Journal of Pain*. 2019(just-accepted).
5. Fletcher D, Stamer UM, Pogatzki-Zahn E, Zaslansky R, Tanase NV, Perruchoud C, et al. Chronic postsurgical pain in Europe: an observational study. *European Journal of Anaesthesiology (EJA)*. 2015;32(10):725-34.
6. Pagé MG, Stinson J, Campbell F, Isaac L, Katz J. Identification of pain-related psychological risk factors for the development and maintenance of pediatric chronic postsurgical pain. *Journal of pain research*. 2013;6:167.
7. Weiser TG, Haynes AB, Molina G, Lipsitz SR, Esquivel MM, Uribe-Leitz T, et al. Size and distribution of the global volume of surgery in 2012. *Bulletin of the World Health Organization*. 2016;94(3):201.
8. Sterling M, Hendrikz J, Kenardy J. Compensation claim lodgement and health outcome developmental trajectories following whiplash injury: a prospective study. *Pain*. 2010;150(1):22-8.
9. Rebbeck T, Leaver A, Bandong A, Kenardy J, Refshauge K, Connelly L, et al. Implementation of a guideline-based clinical pathway of care to improve health outcomes following whiplash injury (Whiplash ImPaCT): protocol of a randomised, controlled trial. *Journal of physiotherapy [Internet]*. 2016; 62(2):[111 p.]
10. Sterling M, Smeets R, Keijzers G, Warren J, Kenardy J. Physiotherapist-delivered stress inoculation training integrated with exercise versus physiotherapy exercise alone for acute whiplash-associated disorder (StressModex): a randomised controlled trial of a combined psychological/physical intervention. *BJSM online*. 2019;19:19.
11. Buchbinder R, van Tulder M, Öberg B, Costa LM, Woolf A, Schoene M, et al. Low back pain: a call for action. *The Lancet*. 2018;391(10137):2384-8.
12. Clark S, Horton R. Low back pain: a major global challenge. *Lancet (London, England)*. 2018;391(10137):2302.
13. Foster NE, Anema JR, Cherkin D, Chou R, Cohen SP, Gross DP, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *The Lancet*. 2018;391(10137):2368-83.
14. Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, et al. What low back pain is and why we need to pay attention. *The Lancet*. 2018;391(10137):2356-67.
15. Vos T, Allen C, Arora M, Barber RM, Bhutta ZA, Brown A, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*. 2016;388(10053):1545-602.
16. Costa LdCM, Maher CG, Hancock MJ, McAuley JH, Herbert RD, Costa LO. The prognosis of acute and persistent low-back pain: a meta-analysis. *Cmaj*. 2012;184(11):E613-E24.
17. Australia SW. Return to Work Survey, 2017/18 Headline Measures Report.: Social Research Centre; 2018.
18. Beales D, Fried K, Nicholas M, Blyth F, Finnis D, Moseley GL. Management of musculoskeletal pain in a compensable environment: Implementation of helpful and unhelpful Models of Care in supporting recovery and return to work. *Best practice & research Clinical rheumatology*. 2016;30(3):445-67.
19. Nicholas M, Costa D, Linton S, Main C, Shaw W, Pearce G, et al. Implementation of Early Intervention Protocol in Australia for 'High Risk'Injured Workers is Associated with Fewer Lost Work Days Over 2 Years Than Usual (Stepped) Care. *J Occup Rehabil*. 2019:1-12.
20. Nicholas M, Costa D, Linton SJ, Main C, Shaw W, Pearce R, et al. Predicting Return to Work in a Heterogeneous Sample of Recently Injured Workers Using the Brief ÖMPSQ-SF. *J Occup Rehabil*. 2019;29(2):295-302.
21. Papagoras H, Pizzari T, Coburn P, Sleigh K, Briggs AM. Supporting return to work through appropriate certification: a systematic approach for Australian primary care. *Australian health review*. 2018;42(2):164-7.

22. Zhang DD, Sussman J, Dossa F, Jivraj N, Ladha K, Brar S, et al. A Systematic Review of Behavioral Interventions to Decrease Opioid Prescribing After Surgery. *Annals of surgery*. 2019.
23. Sun EC, Darnall BD, Baker LC, Mackey S. Incidence of and risk factors for chronic opioid use among opioid-naive patients in the postoperative period. *JAMA internal medicine*. 2016;176(9):1286-93.
24. Katz J, Weinrib A, Fashler SR, Katznelson R, Shah BR, Ladak SS, et al. The Toronto General Hospital Transitional Pain Service: development and implementation of a multidisciplinary program to prevent chronic postsurgical pain. *Journal of pain research*. 2015;8:695.
25. Hsu RJ, Mir KH, Wally BM, Seymour BR. Clinical Practice Guidelines for Pain Management in Acute Musculoskeletal Injury. *Journal of Orthopaedic Trauma*. 2019;33(5):e158-e82.
26. Chou R, Gordon DB, de Leon-Casasola OA, Rosenberg JM, Bickler S, Brennan T, et al. Management of Postoperative Pain: a clinical practice guideline from the American pain society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' committee on regional anesthesia, executive committee, and administrative council. *The Journal of Pain*. 2016;17(2):131-57.
27. Authority MA. Guidelines for the management of acute whiplash-associated disorders for health professionals. Sydney, Australia: Motor Accidents Authority. 2014.
28. Management of people with acute low back pain: model of care. NSW Agency for Clinical Innovation NSW Health; 2016.
29. Services USDoHaH. Pain Management Best Practices Inter-Agency Task Force Report: Updates, Gaps, Inconsistencies, and Recommendations. . May, 2019.
30. Almeida M, Saragiotto B, Richards B, Maher CG. Primary care management of non-specific low back pain: key messages from recent clinical guidelines. *Medical Journal of Australia*. 2018;208(6):272-5.
31. Linton SJ, Nicholas M, Shaw W. Why wait to address high-risk cases of acute low back pain? A comparison of stepped, stratified, and matched care. *Pain*. 2018;159(12):2437-41.
32. Slater H, Dear BF, Merolli MA, Li LC, Briggs AM. Use of eHealth technologies to enable the implementation of musculoskeletal Models of Care: Evidence and practice. *Best Practice & Research Clinical Rheumatology*. 2016;30(3):483-502.
33. van Egmond MA, van der Schaaf M, Vredevelde T, Vollenbroek-Hutten MMR, van Berge Henegouwen MI, Klinkenbijn JHG, et al. Effectiveness of physiotherapy with telerehabilitation in surgical patients: a systematic review and meta-analysis. *Physiotherapy*. 2018;104(3):277-98.
34. Weinrib AZ, Azam MA, Birnie KA, Burns LC, Clarke H, Katz J. The psychology of chronic post-surgical pain: new frontiers in risk factor identification, prevention and management. *Br*. 2017;11(4):169-77.
35. Marin TJ, Van Eerd D, Irvin E, Couban R, Koes BW, Malmivaara A, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain. *Cochrane Database of Systematic Reviews*. 2017(6).
36. Meyer C, Denis CM, Berquin AD. Secondary prevention of chronic musculoskeletal pain: a systematic review of clinical trials. *Annals of physical and rehabilitation medicine*. 2018;61(5):323-38.
37. Monticone M, Cedraschi C, Ambrosini E, Rocca B, Fiorentini R, Restelli M, et al. Cognitive-behavioural treatment for subacute and chronic neck pain. *Cochrane Database of Systematic Reviews*. 2015(5).
38. Nicholls JL, Azam MA, Burns LC, Englesakis M, Sutherland AM, Weinrib AZ, et al. Psychological treatments for the management of postsurgical pain: a systematic review of randomized controlled trials. *Patient related outcome measures*. 2018;9:49.
39. Schröter J, Lechterbeck M, Hartmann F, Gercek E. Structured rehabilitation after lumbar spine surgery : subacute treatment phase. *Der orthopade*. 2014;43(12):1089-95.
40. Watson JA, Ryan CG, Cooper L, Ellington D, Whittle R, Lavender M, et al. Pain neuroscience education for adults with chronic musculoskeletal pain: a mixed-methods systematic review and meta-analysis. *The Journal of Pain*. 2019.
41. Nicholas MK, Linton SJ, Watson PJ, Main CJ, Group DotFW. Early identification and management of psychological risk factors ("yellow flags") in patients with low back pain: a reappraisal. *Physical therapy*. 2011;91(5):737-53.
42. Tawfic Q, Kumar K, Pirani Z, Armstrong K. Prevention of chronic post-surgical pain: the importance of early identification of risk factors. In: Tawfic Q, editor. 2017. p. 424-31.
43. Thapa P, Euasobhon P. Chronic postsurgical pain: current evidence for prevention and management. *Korean J Pain*. 2018;31(3):155-73.

44. Gewandter JS, Dworkin RH, Turk DC, Farrar JT, Fillingim RB, Gilron I, et al. Research design considerations for chronic pain prevention clinical trials: IMMPACT recommendations. *Pain*. 2015;156(7):1184.
45. VanDenKerkhof EG, Peters ML, Bruce J. Chronic pain after surgery: time for standardization? A framework to establish core risk factor and outcome domains for epidemiological studies. *The Clinical journal of pain*. 2013;29(1):2-8.
46. Althaus A, Hinrichs-Rocker A, Chapman R, Becker OA, Lefering R, Simanski C, et al. Development of a risk index for the prediction of chronic post-surgical pain. *European Journal of Pain*. 2012;16(6):901-10.
47. Foster NE, Mullis R, Hill JC, Lewis M, Whitehurst DG, Doyle C, et al. Effect of stratified care for low back pain in family practice (IMPACT Back): a prospective population-based sequential comparison. *The Annals of Family Medicine*. 2014;12(2):102-11.
48. Hill JC, Whitehurst DG, Lewis M, Bryan S, Dunn KM, Foster NE, et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *The Lancet*. 2011;378(9802):1560-71.
49. Ritchie C, Hendrikz J, Kenardy J, Sterling M. Derivation of a clinical prediction rule to identify both chronic moderate/severe disability and full recovery following whiplash injury. *PAIN®*. 2013;154(10):2198-206.
50. Kelly J, Sterling M, Rebbeck T, Bandong AN, Leaver A, Mackey M, et al. Health practitioners' perceptions of adopting clinical prediction rules in the management of musculoskeletal pain: a qualitative study in Australia. *BMJ Open*. 2017;7(8):e015916.
51. Murgatroyd DF, Casey PP, Cameron ID, Harris IA. The effect of financial compensation on health outcomes following musculoskeletal injury: systematic review. *PLoS one*. 2015;10(2):e0117597.
52. Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychological bulletin*. 2007;133(4):581.
53. Bevers K, Watts L, Kishino ND, Gatchel RJ. The biopsychosocial model of the assessment, prevention, and treatment of chronic pain. *US Neurol*. 2016;12(2):98-104.
54. de C Williams AC, Eccleston C, Morley S. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane database of systematic reviews*. 2012(11).
55. Joypaul S, Kelly FS, King MA. Turning Pain into Gain: Evaluation of a Multidisciplinary Chronic Pain Management Program in Primary Care. *Pain Med*. 2018.
56. Keefe FJ, Main CJ, George SZ. Advancing psychologically informed practice for patients with persistent musculoskeletal pain: promise, pitfalls, and solutions. *Physical therapy*. 2018;98(5):398-407.
57. Traeger A, Lee H, Hübscher M, Skinner I, Moseley G, Nicholas M, et al. Effect of Intensive Patient Education vs Placebo Patient Education on Outcomes in Patients With Acute Low Back Pain: A Randomized Clinical Trial. *JAMA Neurology*. 2019;76(2):161.
58. Government A. National Strategic Action Plan for Pain Management. Department of Health; 2019.
59. Lin I, Green C, Bessarab D. 'Yarn with me': applying clinical yarning to improve clinician-patient communication in Aboriginal health care. *Australian Journal of Primary Health*. 2016;22(5):377-82.
60. Ibrahim MS, Khan MA, Nizam I, Haddad FS. Peri-operative interventions producing better functional outcomes and enhanced recovery following total hip and knee arthroplasty: an evidence-based review. *BMC medicine*. 2013;11(1):37.
61. Wiangkham T, Duda J, Haque S, Madi M, Rushton A. The effectiveness of conservative management for acute whiplash associated disorder (WAD) II: a systematic review and meta-analysis of randomised controlled trials. *PLoS One*. 2015;10(7):e0133415.
62. Qaseem A, Wilt TJ, McLean RM, Forciea MA. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Annals of internal medicine*. 2017;166(7):514-30.
63. National GCU. Low Back Pain and Sciatica in Over 16s: Assessment and Management. 2016.
64. Stochkendahl MJ, Kjaer P, Hartvigsen J, Kongsted A, Aaboe J, Andersen M, et al. National Clinical Guidelines for non-surgical treatment of patients with recent onset low back pain or lumbar radiculopathy. *European Spine Journal*. 2018;27(1):60-75.
65. Huijnen IP, Rusu AC, Scholich S, Meloto CB, Diatchenko L. Subgrouping of low back pain patients for targeting treatments: evidence from genetic, psychological, and activity-related behavioral approaches. *The Clinical journal of pain*. 2015;31(2):123-32.

66. Sowden G, Hill JC, Konstantinou K, Khanna M, Main CJ, Salmon P, et al. Targeted treatment in primary care for low back pain: the treatment system and clinical training programmes used in the IMPaCT Back study (ISRCTN 55174281). *Family practice*. 2011;29(1):50-62.
67. Lin I, O'Sullivan P, Coffin J, Mak D, Toussaint S, Straker L. 'I am absolutely shattered': The impact of chronic low back pain on Australian Aboriginal people. *European Journal of Pain*. 2012;16(9):1331-41.
68. Lin IB, O'Sullivan PB, Coffin JA, Mak DB, Toussaint S, Straker LM. Disabling chronic low back pain as an iatrogenic disorder: a qualitative study in Aboriginal Australians. *BMJ Open*. 2013;3(4):e002654.
69. Lin I, O'Sullivan P, Coffin J, Mak D, Toussaint S, Straker L. 'I can sit and talk to her': Aboriginal people, chronic low back pain and healthcare practitioner communication. *Australian family physician*. 2014;43(5):320.
70. Lin IB, Ryder K, Coffin J, Green C, Dalgety E, Scott B, et al. Addressing disparities in low back pain care by developing culturally appropriate information for aboriginal australians: "My Back on Track, My Future". *Pain Med*. 2017;18(11):2070-80.
71. Cullen K, Irvin E, Collie A, Clay F, Gensby U, Jennings P, et al. Effectiveness of workplace interventions in return-to-work for musculoskeletal, pain-related and mental health conditions: an update of the evidence and messages for practitioners. *J Occup Rehabil*. 2018;28(1):1-15.
72. Jellema P, Windt van der, Windt van der, van der Horst HE, Blankenstein AH, Bouter LM, Stalman WAB: Why is a treatment aimed at psychosocial factors not effective in patients with (sub) acute low back pain. *Pain*. 2005;118:350-9.
73. Mathieson S. Patient and clinician interventions to assist deprescribing opioid analgesics in chronic non-cancer pain: a systematic review. *Annals of Family Medicine*. 2019;Under review.
74. Sullivan MD, Turner JA, DiLodovico C, D'Appollonio A, Stephens K, Chan Y-F. Prescription opioid taper support for outpatients with chronic pain: a randomized controlled trial. *The Journal of Pain*. 2017;18(3):308-18.
75. Darnall BD, Ziadni MS, Krishnamurthy P, Flood P, Heathcote L, Mackey IG, et al. "My Surgical Success": Effect of a Digital Behavioral Pain Medicine Intervention on Time to Opioid Cessation After Breast Cancer Surgery-A Pilot Randomized Controlled Clinical Trial. *Pain Med*. 2019;13:13.
76. White R, Hayes C, White S, Hodson FJ. Using social media to challenge unwarranted clinical variation in the treatment of chronic noncancer pain: The "Brainman" story. *Journal of Pain Research*. 2016;9:701-9.
77. Archer KR, Devin CJ, Vanston SW, Koyama T, Phillips SE, George SZ, et al. Cognitive-behavioral-based physical therapy for patients with chronic pain undergoing lumbar spine surgery: a randomized controlled trial. *The Journal of Pain*. 2016;17(1):76-89.
78. Hesselink G, Schoonhoven L, Barach P, Spijker A, Gademan P, Kalkman C, et al. Improving patient handovers from hospital to primary care. *Ann Intern Med*. 2012;157(6):417-28.
79. Ensing HT, Stuijt CC, Van Den Bemt BJ, Van Dooren AA, Karapinar-Carkit F, Koster ES, et al. Identifying the optimal role for pharmacists in care transitions: a systematic review. *Journal of managed care & specialty pharmacy*. 2015;21(8):614-36.
80. Bethishou L, Herzik K, Fang N, Abdo C, Tomaszewski DM. The impact of the pharmacist on continuity of care during transitions of care: A systematic review. *Journal of the American Pharmacists Association*. 2019.
81. Chronic Pain Telehealth Pilot Project Evaluation Report. NSW Agency for Clinical Innovation; 2016.
82. Improving physiotherapy access using telehealth. Murrumbidgee Local Health District. NSW Agency for Clinical Innovation 2018.
83. Gentry MT, Lapid MI, Clark MM, Rummans TA. Evidence for telehealth group-based treatment: A systematic review. *Journal of telemedicine and telecare*. 2019;25(6):327-42.
84. Mariano TY, Wan L, Edwards RR, Lazaridou A, Ross EL, Jamison RN. Online group pain management for chronic pain: Preliminary results of a novel treatment approach to teletherapy. *Journal of telemedicine and telecare*. 2019;1357633X19870369.
85. Telehealth in practice guide NSW Agency for Clinical Innovation 2019. Report No.: ISBN: 978-1-76081-282-9 (print) ; 978-1-76081-283-6 (online); (ACI) 190551.
86. Dear BF, Gandy M, Karin E, Fogliati R, Fogliati VJ, Staples LG, et al. The Pain Course: 12- and 24-Month Outcomes From a Randomized Controlled Trial of an Internet-Delivered Pain Management Program Provided With Different Levels of Clinician Support. *The Journal of Pain*. 2018;19(12):1491-503.

87. Dear BF, Gandy M, Karin E, Staples LG, Johnston L, Fogliati VJ, et al. The Pain Course: a randomised controlled trial examining an internet-delivered pain management program when provided with different levels of clinician support. *Pain*. 2015;156(10):1920.
88. Schultz R, Smith J, Newby JM, Gardner T, Shiner CT, Andrews G, et al. Pilot Trial of the Reboot Online Program: An Internet-Delivered, Multidisciplinary Pain Management Program for Chronic Pain. *Pain Research and Management*. 2018;2018.
89. Reynoldson C, Stones C, Allsop M, Gardner P, Bennett MI, Closs SJ, et al. Assessing the quality and usability of smartphone apps for pain self-management. *Pain Med*. 2014;15(6):898-909.
90. McKay FH, Cheng C, Wright A, Shill J, Stephens H, Uccellini M. Evaluating mobile phone applications for health behaviour change: a systematic review. *Journal of telemedicine and telecare*. 2018;24(1):22-30.
91. Machado GC, Pinheiro MB, Lee H, Ahmed OH, Hendrick P, Williams C, et al. Smartphone apps for the self-management of low back pain: A systematic review. *Best Practice & Research Clinical Rheumatology*. 2016;30(6):1098-109.
92. Hall A, Richmond H, Copsey B, Hansen Z, Williamson E, Jones G, et al. Physiotherapist-delivered cognitive-behavioural interventions are effective for low back pain, but can they be replicated in clinical practice? A systematic review. *Disability and rehabilitation*. 2018;40(1):1-9.
93. Beales D, McManus L, Tan J-S, Elliott C, Mitchell T. Implementation of Questionnaire-Based Risk Profiling for Clients in a Workers' Compensation Environment: An Example in Australian Physiotherapy Practice. *J Occup Rehabil*. 2019:1-8.
94. Daly A. Worklessness: can physiotherapists do more? *Journal of physiotherapy*. 2016;62(4):179-80.
95. Synnott A, O'Keeffe M, Bunzli S, Dankaerts W, O'Sullivan P, O'Sullivan K. Physiotherapists may stigmatise or feel unprepared to treat people with low back pain and psychosocial factors that influence recovery: a systematic review. *Journal of physiotherapy*. 2015;61(2):68-76.
96. Cowell I, O'Sullivan P, O'Sullivan K, Poyton R, McGregor A, Murtagh G. The perspectives of physiotherapists on managing nonspecific low back pain following a training programme in cognitive functional therapy: A qualitative study. *Musculoskelet*. 2019;17(1):79-90.
97. Cowell I, O'Sullivan P, O'Sullivan K, Poyton R, McGregor A, Murtagh G. Perceptions of physiotherapists towards the management of non-specific chronic low back pain from a biopsychosocial perspective: A qualitative study. *Musculoskeletal Science and Practice*. 2018;38:113-9.
98. Kelly JM, Bunzli S, Ritchie C, Kenardy J, Smeets R, Sterling M. Physiotherapist-delivered Stress Inoculation Training for acute whiplash-associated disorders: A qualitative study of perceptions and experiences. *Musculoskeletal science and practice*. 2018;38:30-6.
99. Arora S, Kalishman SG, Thornton KA, Komaromy MS, Katzman JG, Struminger BB, et al. Project ECHO: A telementoring network model for continuing professional development. *J Contin Educ Health Prof*. 2017;37(4):239-44.
100. Serhal E, Arena A, Sockalingam S, Mohri L, Crawford A. Adapting the consolidated framework for implementation research to create organizational readiness and implementation tools for project ECHO. *The Journal of continuing education in the health professions*. 2018;38(2):145.
101. Katzman JG, Galloway K, Olivas C, McCoy-Stafford K, Duhigg D, Comerchi G, et al. Expanding health care access through education: dissemination and implementation of the ECHO model. *Military medicine*. 2016;181(3):227-35.
102. Shelley BM, Katzman JG, Comerchi Jr GD, Duhigg DJ, Olivas C, Kalishman S, et al. ECHO pain curriculum: Balancing mandated continuing education with the needs of rural health care practitioners. *J Contin Educ Health Prof*. 2017;37(3):190-4.
103. Frank JW, Carey EP, Fagan KM, Aron DC, Todd-Stenberg J, Moore BA, et al. Evaluation of a telementoring intervention for pain management in the Veterans Health Administration. *Pain Med*. 2015;16(6):1090-100.
104. Furlan AD, Zhao J, Voth J, Hassan S, Dubin R, Stinson JN, et al. Evaluation of an innovative tele-education intervention in chronic pain management for primary care clinicians practicing in underserved areas. *Journal of telemedicine and telecare*. 2018:1357633X18782090.
105. Ball S, Wilson B, Ober S, Mchaourab A. SCAN-ECHO for pain management: Implementing a regional telementoring training for primary care providers. *Pain Med*. 2017;19(2):262-8.

106. Carlin L, Zhao J, Dubin R, Taenzer P, Sidrak H, Furlan A. Project ECHO telementoring intervention for managing chronic pain in primary care: Insights from a qualitative study. *Pain Med.* 2017;19(6):1140-6.
107. Zhou C, Crawford A, Serhal E, Kurdyak P, Sockalingam S. The impact of project ECHO on participant and patient outcomes: a systematic review. *Academic Medicine.* 2016;91(10):1439-61.
108. Shimasaki S, Bishop E, Guthrie M, Thomas JF. Strengthening the Health Workforce through the ECHO Stages of Participation: Participants' Perspectives on Key Facilitators and Barriers. *Journal of medical education and curricular development.* 2019;6:2382120518820922.
109. Stanley B, Jackson A, Norman A, Collins L, Bonomo A, Bonomo Y. Opioid prescribing improvement in orthopaedic specialty unit in a tertiary hospital: a retrospective audit of hospital discharge data pre-and post-intervention for better opioid prescribing practice. *ANZ J Surg.* 2019.
110. Holliday S, Hayes C, Dunlop A, Morgan S, Tapley A, Henderson K, et al. Protecting Pain Patients. The Evaluation of a Chronic Pain Educational Intervention. *Pain Med.* 2017;18(12):2306-15.
111. Gray JS, Swan JR, Lynch MA, Tay TM, Mackenzie MJ, Wiggers JH, et al. Hunter and New England HealthPathways: a 4-year journey of integrated care. *Aust Health Rev.* 2018;42(1):66-71.
112. Norris S, Schneider CH, Applegarth K, Wortley S, Elshaug A, Wilson A. Health Pathways Sydney Evaluation: Final Report. 2018.
113. Stokes T, Tumilty E, Doolan-Noble F, Gauld R. HealthPathways implementation in a New Zealand health region: a qualitative study using the Consolidated Framework for Implementation Research. *BMJ Open.* 2018;8(12):e025094.
114. Gill SD, Mansfield S, McLeod M, Von Treuer K, Dunn M, Quirk F. HealthPathways improving access to care. *Australian Health Review.* 2019;43(2):207-16.
115. Tardif H, Arnold C, Hayes C, Eagar K. Establishment of the Australasian electronic persistent pain outcomes collaboration. *Pain Med.* 2016;18(6):1007-18.
116. Brunner E, De Herdt A, Minguet P, Baldew S-S, Probst M. Can cognitive behavioural therapy based strategies be integrated into physiotherapy for the prevention of chronic low back pain? A systematic review. *Disability and rehabilitation.* 2013;35(1):1-10.
117. Sullivan MJ, Adams H, Rhodenizer T, Stanish WD. A psychosocial risk factor-targeted intervention for the prevention of chronic pain and disability following whiplash injury. *Physical therapy.* 2006;86(1):8-18.
118. Katzman JG, Qualls CR, Satterfield WA, Kistin M, Hofmann K, Greenberg N, et al. Army and Navy ECHO Pain Telementoring Improves Clinician Opioid Prescribing for Military Patients: an Observational Cohort Study. *Journal of general internal medicine.* 2019;34(3):387-95.
119. Katzman JG, Comerchi Jr G, Boyle JF, Duhigg D, Shelley B, Olivas C, et al. Innovative telementoring for pain management: project ECHO pain. *J Contin Educ Health Prof.* 2014;34(1):68-75.
120. Flynn DM, Eaton LH, McQuinn H, Alden A, Meins AR, Rue T, et al. TelePain: Primary care chronic pain management through weekly didactic and case-based telementoring. *Contemporary clinical trials communications.* 2017;8:162-6.

Appendix 1: Chronic Pain Project Steering Committee

Ms Carol Bennett, CEO, painaustralia

Mr David Beveridge, Nurse Practitioner, Lismore Base Hospital, Multidisciplinary Pain Management Clinic

Dr Matthew Bryant, Director Townsville Pain Persistent Pain Service and NQPPMS

Sr Mary-Lynne Cochrane, Consumer Representative

Dr Anne Daly, Physiotherapy and Pain Management Consultant

Ms Terina Grace, CEO and Managing Director Black Swan Health

Ms Fiona Hodson, Clinical Nurse Consultant Pain Management, Hunter Integrated Pain Service, Surgical Services

Associate Professor Malcolm Hogg, painaustralia

Dr Simon Holliday, GP and Addiction Medicine Specialist

Ms Jenni Johnson, Manager, Pain Management Network, NSW ACI (February 2018-June 2019)

Ms Susan Rogers, Manager, Pain Management Network, NSW ACI (July 2019-)

Ms Margaret Knight, Consumer Representative

Ms Joyce McSwan, Pharmacist, Pain Educator Gold Coast PHN

Professor Michael Nicholas, Director, Pain Education & Pain Management Programs, PMRI, University of Sydney

Dr Milana Votrubic, GP specialising in pain

Ms Leanne Wells, Consumers Health Forum and consumer representative on Pain Australia

Professor Andrew Wilson, Director, TAPPC and Co-Director Menzies Centre for Health Policy

Appendix 2: Scope of the rapid review

Purpose of the review

The purpose of the rapid evidence review was to inform a deliberative dialogue with PHNs about the secondary prevention of chronic pain.

Research questions

The rapid review sought to answer the following questions:

- i. What are the key principles related to the secondary prevention of chronic pain highlighted in the evidence?
- ii. What strategies related to Goal 1 (consumer and community initiatives) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)? Six key focus areas include:
 - a. pain that is multifocal or not condition specific
 - b. surgery
 - c. whiplash
 - d. low back pain
 - e. return to work and work-related injuries
 - f. opioid-related initiatives
- iii. What strategies related to Goal 2 (health professional capacity building) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)?
- iv. What strategies related to Goal 3 (health system support) have been shown to be effective for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)?

The rapid evidence review did not aim to systematically search for, or synthesise, all the relevant evidence and did not aim to provide an indication of the strength of the evidence in the selected bodies of evidence. The rapid evidence review provides a narrative synthesis of the evidence and highlights case study examples. The evidence informs the development of the *Principles for the secondary prevention of chronic pain* and the mapping of the options for PHNs to address the secondary prevention and management of chronic pain.

Validation of the rapid evidence review

The scoping of the rapid evidence review, development of the search strategy, conducting of the literature search and synthesising the results was undertaken by the first author (SDM). Another author (LM) conducted the grey literature search with input by the first author (SDM). The rapid evidence review was reviewed by key stakeholders and the research team to check for any gaps. The final draft of the rapid evidence review was reviewed by the other authors (FB, LM, MM, DS and MM) and feedback incorporated prior to being finalised and sent to participants to inform the PHN deliberative dialogue.

Search strategy

The following search strategy was used in the rapid evidence review:

1. Medline, PsychINFO, Cochrane Database of Systematic Reviews, Joanna Briggs Institute EBP Database and PEDro database search of the peer-review literature from Australia, UK, Europe, USA, Canada and New Zealand, 2010-2019 (in the English language)

2. Hand searching of references from relevant papers
3. Google and Google Scholar search of the grey literature from Australia and internationally
4. Grey literature search of key agencies from Australia and internationally to identify evaluation reports, initiatives (without evaluations) and health professional and consumer resources
5. Relevant peer-review literature, evaluation reports and initiatives (without evaluations) identified in the consultation with PHNs in Phase 1 of the Chronic Pain Project
6. Key stakeholders (with content expertise in pain management and related research)⁷ were asked to identify relevant peer-review literature, evaluation reports, initiatives (without evaluations), health professional and consumer resources, and research studies and initiatives 'in the pipeline'.
7. The final evidence review was reviewed by key stakeholders and the research team to check for any gaps.

Inclusion criteria

1. Peer-review publications, evaluation reports and clinical practice guidelines from Australia, UK, Europe, USA, Canada and New Zealand, 2010-2019 (in the English language)
2. Study designs include:
 - a. Clinical practice guidelines
 - b. Narrative reviews and scoping reviews
 - c. Systematic reviews
 - d. Experimental or quasi-experimental studies or peer-review protocols
 - e. Pre-post-test studies, post-test studies (observational studies)
3. Studies relate to consumer and community initiatives for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain) in the six key focus areas (see above)
4. Studies relate to health professional capacity building for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)
5. Studies relate to health system support for the secondary prevention of chronic pain (or the management of chronic pain with the potential for adaptation to the secondary prevention of chronic pain)
6. Studies implemented in the primary care setting
7. Studies implemented in hospital and compensable settings that have the potential to be adapted to the primary care setting.

Exclusion criteria

1. Studies relate to cancer pain

⁷ Professor Helen Slater, Dr Duncan Sanders, Ms Fiona Hodson, Professor Michael Nicholas, Dr Simon Holliday, Ms Joyce McSwan, Dr Stephanie Mathieson

2. Studies relate to consumer initiatives about pharmacotherapy; or surgical techniques; or physical therapy techniques

Selection of the case study examples

Case studies examples were selected to demonstrate recent interventions related to the secondary prevention and management of chronic pain and included a range of healthcare providers and modes (e.g. face-to-face, online). Case study examples from the primary care setting were prioritised in this review. Case study examples from hospital and compensable settings were included in the review in areas where there was a lack of evidence relating to interventions in the primary care setting.

Grey literature search

1. Grey literature search of key agencies and agencies listed on International Association for the Study of Pain (IASP) to identify evaluation reports, initiatives (without evaluations) and health professional and consumer resources. A list of agencies searched is below.
2. Google advanced search technique was used to conduct 3 searches of 1) all Australian organisation websites (Search 1 = "site:org.au") 2) all Australian government websites (Search 2 = "site:gov.au") and 3) educational institution websites (Search 3 = "site:edu"), each containing key search terms, including
 - i. acute pain OR subacute pain OR sub-acute pain AND prevention OR preventative health services OR early intervention OR secondary prevention
3. Ad hoc identification of literature, through internet searching, full text readings, reference list scanning or documents that have been identified through stakeholders.

The following websites were searched for relevant grey literature.

Pain-related agencies

- Painaustralia <https://www.painaustralia.org.au>
- Australian Pain Society <https://www.apsoc.org.au/>
- Australian Pain Management Association <https://www.painmanagement.org.au/>
- Chronic Pain Australia <http://chronicpainaustralia.org.au/>
- Arthritis Australia <https://arthritisaustralia.com.au/> and <https://mybackpain.org.au/>
- Musculoskeletal Australia <https://www.msk.org.au/>
- International Association for the Study of Pain (IASP) <https://www.iasp-pain.org/>
- American Chronic Pain Association <http://www.theacpa.org>
- Australian Pain Management Association <http://www.painmanagement.org.au/>
- Canadian Pain Coalition <http://www.canadianpaincoalition.ca/>
- Chronic Pain Australia <http://www.chronicpainaustralia.org.au/>
- Patient Advocate Foundation <http://www.patientadvocate.org>
- Chronic Pain Scotland <http://chronicpainscotland.org/>
- Pain Connection-Chronic Pain Outreach Center <http://www.painconnection.org/>
- Pain Health WA <https://painhealth.csse.uwa.edu.au/>
- Scottish Society of Acute Pain Services <https://www.ssaps.scot.nhs.uk/i>
- The British Pain Society <https://www.britishpainsociety.org/>

Other agencies

- Departments of Health (Federal, WA, SA, NT, QLD, NSW, Vic, ACT, Tas)
- NSW Agency for Clinical Innovation (ACI) <https://www.aci.health.nsw.gov.au/>
- National Prescribing Service (NPS) MedicineWise <https://www.nps.org.au/>
- Sax Institute <https://www.saxinstitute.org.au/>

- Consumers Health Forum of Australia <https://chf.org.au/>
- Therapeutic Goods Administration (TGA) <https://www.tga.gov.au/>
- HealthPathways - <https://www.swsphn.com.au/healthpathways>

Health professional associations

- Australian College of Rural and Remote Medicine <https://www.acrrm.org.au/>
- Pharmaceutical Society of Australia <https://www.psa.org.au/>
- Australian Psychology Society <https://www.psychology.org.au/>
- Royal Australian College of Physicians (RACGP) <https://www.racgp.org.au/>
- Australian and New Zealand College of Anaesthetists <http://www.anzca.edu.au/>
- Australian Physiotherapy Association <https://australian.physio/>
- The Australian Clinical Psychology Association <https://acpa.org.au/>

Limitations of the review

This rapid evidence review for PHN decision-makers aimed to maintain methodological rigour while meeting the needs and timelines of the end-user. The review was intended to provide a narrative synthesis of the evidence and highlight case study examples to inform the mapping of the options for PHNs and did not include a grade strength of evidence assessment.

Appendix 3: Evidence map of the options related to Goal 1

Table 2: Evidence map of the options related to Goal 1

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Face-to-face multidisciplinary consumer pain program - for consumers at risk of developing chronic pain	<p>Meyer 2018 systematic review- secondary prevention of (sub)acute pain(36)</p> <p>Marin 2017 systematic review - Multidisciplinary biopsychosocial rehabilitation - subacute low back pain(35)</p> <p>Katz 2015- Toronto Transitional Pain Service - post surgery(24)</p>	Joypaul 2018 - chronic pain)(55)		Face-to-face multidisciplinary consumer pain programs currently implemented in six PHNs and WAPHA.			Potential prototype of a consumer pain program for people with (sub)acute pain who are at risk of developing chronic pain, based on the Turning Pain Into Gain program (see Appendix 9)

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
<p>Psychologically informed physical therapy program provided by a physiotherapist - for consumers at risk of developing chronic pain</p>	<p>Brunner 2013 systematic review- cognitive behaviour therapy-based treatments - (sub)acute low back pain(116)</p> <p>Hall 2018 systematic review - physiotherapist-led cognitive-behavioural interventions - low back pain(92)</p> <p>Nicholls 2018 systematic review – perioperative cognitive behavioural therapy(38)</p> <p>Foster 2014 (IMPACT Back) --- stratified care-psychologically informed physical therapy - (sub)acute and chronic pain(47)</p> <p>Sterling 2019 (StressModex) - physiotherapist-led intervention of stress inoculation training</p>	<p>Foster (IMPACT Back) 2014 – (sub)acute and chronic(47)</p>					

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
<p>Continued.....</p> <p>Psychologically informed physical therapy program provided by a physiotherapist - for consumers at risk of developing chronic pain</p>	<p>and exercise - acute whiplash(10)</p> <p>Sullivan 2006 -psychological intervention and physical therapy in whiplash(117)</p> <p>Archer 2016 - cognitive-behavioural based physical therapy – post-surgery(77)</p> <p>Nicholas 2019 -early intervention for injured workers- compensation environment(19)</p>						
<p>A consumer education and/or behavioural initiative about safe and effective use of</p>	<p>Zang 2019- systematic review post surgery (hospital setting)(22)</p> <p>Darnall 2019- breast cancer surgery- digital behavioural intervention(75)</p>	<p>Mathieson 2019 systematic review -primary care setting (under review)(73)</p> <p>Sullivan 2017(74)</p>	<p>Opioid Early Intervention Pilot Project (West Vic PHN)</p>				

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
medications and how to taper opioids		Holliday 2017 (110) White 2016- Brainman video series(76)					
Transitions of care / pre-surgery consumer initiative	Chou 2016-Clinical guidelines-pain(26) Shug 2015 -Acute pain management scientific evidence(3) Nicholls 2018- psychological treatments-surgery(38) Darnall 2019- breast cancer surgery- digital behavioural intervention(75) Archer 2016 – cognitive behavioural therapy- lumbar spine surgery(77)						

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Transitions of care / post-surgery consumer initiative	<p>Hesselink 2012 - systematic review- to improve patient handovers from hospital to primary care(78)</p> <p>Ensing 2015 systematic review- community pharmacists(79)</p> <p>Bethishou 2019 systematic review- community pharmacists(80)</p>		<p>The Pain Prescribing on Discharge Working Group including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health.</p>				
Telehealth assisted allied health services (individual/group sessions)	<p>Egmond 2018 systematic review – physiotherapy with telerehabilitation(33)</p>	<p>Mariano 2019- group based program- chronic pain(84)</p>		<p>Outreach services (telehealth and visiting health care providers) connecting people</p>		<p>The NSW ACI. Improving physiotherapy access using telehealth.</p>	

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Continued.... Telehealth assisted allied health services (individual/group sessions)	Gentry 2019- telehealth group-based programs (range of conditions)(83)			in pain with pain specialists and other health providers		Murrumbidgee Local Health District, Report, 2018(82) NSW ACI. Chronic Pain -Telehealth Pilot Project Evaluation Report 2016(81)	
Community awareness campaign		White 2016- Brainman video series(76)	Support for Pain Revolution Brainman video series (could be further adapted)				
Face-to-face and/or online peer support group (network)				Face-to-face peer support network - Adelaide PHN			

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Online consumer pain program		Dear 2018, 2015 - The Pain Course (86, 87) Schultz 2018 - This Way Up (88)					
Mobile app for the post-surgery or post-injury phase or (sub)acute back pain		Machado 2016- systematic review- low back pain (91) Reynoldson 2014 review – pain self-management (89) McKay 2018 systematic review - health behaviour					

Goal 1: Access to multidisciplinary care and improving consumer health literacy and care navigation

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
		change (range of conditions) (90)					
Promotion of relevant consumer resources	N/A	N/A	Promotion of consumer resources: <ul style="list-style-type: none"> ➤ Via consumer and health professional networks (events and newsletters), ➤ Via HealthPathways ➤ Via online consumer distribution platforms & information portals implemented in four PHNs (e.g. GoShare, Health Resource Directory, Patient Info) 		N/A	N/A	

Appendix 4: Evidence map of the options related to Goal 2

Table 3: Evidence map of the options related to Goal 2

Goal 2: Ensuring health professionals are skilled and provide best-practice evidence-based care							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Face-to-face and/or online education and training for primary care providers (GPs, practice nurses, allied health practitioners, community pharmacists)	<p>Sowden 2012 (IMPaCT Back Pain) training programme(66)</p> <p>Kelly 2018 (StressModex) – acute whiplash-stress inoculation training(98)</p> <p>Brunner 2013 Systematic review (sub)acute low back pain(116)</p> <p>Beales 2019- implementation of screening tool – physiotherapy</p>	<p>Sowden 2012 (IMPaCT Back Pain) training programme(66)</p> <p>Keefe 2018 systematic review - psychologically informed practice for pain management(56)</p> <p>Cowell 2019 – cognitive functional therapy(96)</p> <p>Furlan 2019(104); Frank 2015(103); Katzman 2019(118), 2014(119); Shelley 2017(102); Flynn 2017(120) (Project ECHO- chronic pain)</p>		<ul style="list-style-type: none"> • Most PHNs provide chronic pain management education events • As part of the consumer pain program in South Eastern NSW PHN, ACI supported facilitators to access webinar skills training in pain management (PMRI) 			

Goal 2: Ensuring health professionals are skilled and provide best-practice evidence-based care

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
<p>Continued...</p> <p>Face-to-face and/or online education and training for primary care providers (GPs, practice nurses, allied health practitioners, community pharmacists)</p>	<p>clinics -return to work(93)</p> <p>Papapagorus 2018 - Return to work-flowchart for certification – primary care(21)</p> <p>Kelly 2017 – clinical prediction rule – acute whiplash(50)</p> <p>Hall 2018- Physiotherapist-delivered cognitive-behavioural interventions- low back pain(92)</p>						
<p>Initiative about opioids</p>	<p>Zhang 2019 systematic review Behavioural Interventions to</p>	<p>Holliday -GPs trainees(110)</p> <p>Mathieson 2019 systematic review -</p>		<p><i>Prescribed Drugs of Dependence Active Learning Modules -</i></p>			

Goal 2: Ensuring health professionals are skilled and provide best-practice evidence-based care

	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Continued... Initiative about opioids	Decrease Opioid Prescribing After Surgery(22) Stanley 2019 - Australian hospital opioid prescribing initiative(109)	primary care setting (under review)		Western Victoria PHN Some PHNs provide education and training events about opioids or embedded in chronic pain management events			
Face-to-face and/or online interdisciplinary community of practice				One PHN (Gold Coast) and WAPHA provide a network for primary care providers involved in a face-to-face consumer pain			
Promotion of relevant education and training and other health professional	N/A	N/A	Some PHNs promote Better Pain Management online education program; webinar skills training in pain management- putting cognitive behavioural therapy skills into practice		N/A	N/A	

Goal 2: Ensuring health professionals are skilled and provide best-practice evidence-based care							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
resources offered by other agencies			(Pain Management Research Institute) NPS MedicineWise educational visits; and Pain Revolution Local Pain Education (LPE) Program				
Other health professional capacity building initiative							

Appendix 5: Evidence map of the options related to Goal 3

Table 4: Evidence map of the options related to Goal 3

Goal 3: Quality improvement and health system support							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
HealthPathways (or other referral pathway system)	<p>Gill 2019- South-West Victoria HealthPathways (not pain specific)(114)</p> <p>Stokes 2018 - New Zealand HealthPathways (not pain specific)(113)</p> <p>Gray 2018 Hunter and New England HealthPathways (not pain specific)(111)</p>		<p>HealthPathways - Most PHNs have implemented HealthPathways</p> <p>Some PHNs have developed relevant referral pathways to assess and manage patients with (sub)acute pain (including post-surgery, post-injury and (sub)acute low back pain populations)</p>		Norris 2018 Sydney HealthPathways (not pain specific)(112)		
Transitional care/discharge planning initiative	Hesselink 2012 - systematic review- to improve patient handovers from hospital to primary care(78)		The Pain Prescribing on Discharge Working Group including the Top End Health Service (TEHS) and the NT PHN, funded by the NT Department of Health.				

Goal 3: Quality improvement and health system support							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Continued... Transitional care/discharge planning initiative	Ensing 2015 systematic review- community pharmacists(79) Bethishou 2019 systematic review- community pharmacists(80)						
Electronic Persistent Pain Outcomes Collaboration (ePPOC)		Tardiff 2016 (115)				ePPOC reports for individual PHNs and pain services	
Prescription drug monitoring systems, e.g. SafeScript			The Victoria PHNs led by Western Victoria PHN have been commissioned to provide education and training (in partnership with NPS MedicineWise) for GPs and pharmacists to support the implementation of SafeScript.				

Goal 3: Quality improvement and health system support							
	Reviews and examples of initiatives from the peer-review literature related to the secondary prevention of chronic pain	Reviews and examples of initiatives from the peer-review literature related to the management of chronic pain	PHN initiatives related to the secondary prevention of chronic pain (PHN consultation Phase 1)	PHN initiatives related to the management of chronic pain (PHN consultation Phase 1)	Examples of initiatives from the grey literature (evaluation reports) related to the secondary prevention of chronic pain	Examples of initiatives from the grey literature (evaluation reports) related to the management of chronic pain	Initiatives 'in the pipeline' related to the secondary prevention of chronic pain
Other health system support initiative			Quality Improvement Practice Incentives Program (QIPIP)				

Appendix 6: Enablers to implementing consumer pain programs identified in Phase 1 of the Chronic Pain Project

Evidence of benefit	<ul style="list-style-type: none"> • Evidence of benefit (that is, initiative shown to be feasible, acceptable and effective through program monitoring and evaluation)
Implemented by other PHNs	<ul style="list-style-type: none"> • Delivered or commissioned by other PHNs and ease of adaptation to the local context
Champions	<ul style="list-style-type: none"> • Clinical local champions (for example, GPs with a special interest) and non-clinical local champions (for example, consumers, managers, administrators, funders)
Working groups	<ul style="list-style-type: none"> • Establishment of a working group with a range of stakeholders to help plan, implement and monitor the initiative (for example, stakeholders from the PHN, hospital pain services, commissioned providers, other funders, consumers)
Standardised processes	<ul style="list-style-type: none"> • Standardised processes for communication and referrals
Health professional networks	<ul style="list-style-type: none"> • Establishment of health professional networks to support the implementation of consumer pain programs
Feedback	<ul style="list-style-type: none"> • Regular feedback from consumers, health professionals and commissioned providers
Promotion and engagement	<ul style="list-style-type: none"> • Promotion of the initiative among general practitioners and engagement of consumers
Monitoring and evaluation	<ul style="list-style-type: none"> • Continuous improvement and adaptation; establishing key indicators to evaluate impact/outcomes for commissioned providers; using standardised data collection systems, for example, ePPOC data collection; and partnership with a university to undertake the evaluation
Adequate resources	<ul style="list-style-type: none"> • Adequate funding and staff to deliver or commission the initiative

Appendix 7: Enablers to implementing education and training initiatives identified in Phase 1 of the Chronic Pain Project

<p>Face-to-face educational events implemented or commissioned by PHNs</p>	<ul style="list-style-type: none"> •Local pain specialist or a GP with a special interest in pain to conduct the education event •Selecting topics of interest, usually based on GP surveys (for example, pain management strategies, opioid management) or responding to policy changes, for example, up-scheduling of codeine •Promoting education events through health professional networks and newsletters •Events accredited by the Royal Australian College of General Practitioners (RACGP) •Running events free of charge and at times that are feasible for primary care providers (usually evening seminars)
<p>Support for implementation of education and training conducted by other agencies e.g. NPS Medicine Wise practice education, Pain Revolution Local Pain Education Program (LPEP)</p>	<ul style="list-style-type: none"> •Practice visits to promote the program increased program enrolments •GP incentives •Maintaining relationships with NPS MedicineWise Clinical Service Specialists through the transition from Medicare Local to Primary Health Network •Champions within the PHN, for example, GP liaison officer who has completed the LPEP in Murrumbidgee
<p>Promotion of webinar training</p>	<ul style="list-style-type: none"> •Funding from NSW ACI for facilitators to access the webinar free of charge
<p>Support for mentorship of primary care providers by pain specialists and other members of the pain service</p>	<ul style="list-style-type: none"> •Time and commitment from pain specialists and other members of the pain service
<p>Telehealth and other online services connecting primary care providers with pain specialists and other health providers</p>	<ul style="list-style-type: none"> •Project ECHO provided free of charge to providers online via Zoom. •Project ECHO is an established program, with training provided in the US.

Appendix 8: Enablers to implementing HealthPathways identified in in Phase 1 of the Chronic Pain Project

Executive level support (local hospital networks and PHNs)

Engagement with local hospital networks to enable specialist involvement

Formal partnerships (working groups) between primary care providers and specialists to develop the referral pathways

Involvement of clinical editors in the the development of the referral pathways

Promotion of HealthPathways and how to use it
(e.g. PHN education events with primary care providers; HealthPathways staff conducting site visits to GP practices; promoting HealthPathways through peak GP bodies such as GP Synergy, Hunter Postgraduate Medical Institute)

Responding to feedback from clinicians and addressing any use or content-related issues

Adequate capacity (staff and time) to develop new referral pathways and update the content and design of new referral pathways as needed

Monitoring of usage of localised referral pathways

Evaluation of HealthPathways

Appendix 9: Potential prototype of a consumer pain program for people with (sub)acute pain who are at risk of developing chronic pain

This project plan is a draft only and it is included in this review as a potential prototype only (yet to be funded and tested). It demonstrates how a consumer pain program for chronic pain (Turning Pain Into Gain) could be adapted to a program for consumers with (sub)acute pain at risk of developing chronic pain.

Permission to publish the project plan below has been provided by Ms Joyce McSwan.

Project Plan

A Project Plan is used to elaborate on the information submitted as part of the Concept Brief which was created in the Initiation phase. In the Planning phase, the contents of the Brief are extended and refined into the Project Plan, after which the Concept Brief is no longer maintained.

Project Name: Turning Pain into Gain	Project Officer: Joyce McSwan – PainWISE Pty Ltd
---	---

Early Intervention self-management program for acute and subacute pain conditions within primary health care	
---	--

Date Submitted:	
------------------------	--

Project Overview

Background

- Early stages of acute/subacute pain are different to chronic pain because patients at this phase of the pain condition are not at the stage of accepting a potential life of pain and associated losses. For many acute/subacute pain sufferers, they are focused on 'getting better' which is a very different focus to chronic pain management where potentially a life long journey of persistent pain will require them to learn management skills as well as managing expectations to enable this. As this is very different to a chronic pain scenario it should be treated through a health pathway dedicated to the unique needs of the early intervention of acute/subacute pain.
- Chronic pain and mental health problems, particularly depression, commonly occur together. In Australia and New Zealand, 40.5% of pain patients captured in ePPOC (Electronic Persistent Pain Outcomes Collaboration) data in 2016 reported also suffering depression and/or anxiety (Ref: Blanchard M, Tardif H, Fenwick N, Blissett C and Eagar K (2017) *Electronic Persistent Pain Outcomes Collaboration Annual Data Report 2016. Australian Health Services Research Institute, University of Wollongong.*)
- Currently approximately 15% of patients referred to the TPIG program are between 3-6 months of a pain condition diagnosis
- Approximately 70% of patients who have been referred who are within 3-6 months of a pain condition diagnosis are discharged by 6-7 months of program participation. At the moment they have to stay a minimum of 6 months even if they are ready to leave earlier as our group based Self-Management Education program is a minimum of 6 months in duration.
- A program stream dedicated to acute/subacute pain management will have 3 main goals: a) faster turn-over of patients which will result in better efficiency of fund utilisation within an acute/subacute pain program; b) allow for the TPIG funding to be dedicated to chronic pain sufferers; and c) prevention of progression to chronic/persistent pain.
- Early education and awareness of the risks of opioids have resulted in reduction of opioid requirements and cessation plan. Approximately 80% of all acute/subacute pain management patients reduced their opioids between 6-7 months from time of commencement during participation within the current TPIG program.
- Current research by the Australian Prevention Partnership Centre have recommended primary health care identifies and manage acute/subacute care patients to reduce the healthcare burden of chronic pain.
- This will be the first pilot study of its kind in primary healthcare in Australia.

Psychosocial factors

Predominantly patient risk for developing chronic pain is predicted better by psychosocial factors than biological factors. Some of the more common psychosocial risk factors are:

- The patient's attitudes and beliefs, emotions, behaviours, family, and workplace.
- The behaviour of health professionals can also have a major influence.
- Beliefs or judgements about pain and injury, how disabling it is.
- Poor work history, or unsupportive work environment.
- Thinking the worst, that the pain is uncontrollable or will never improve.
- Comorbid depression, anxiety, or social withdrawal.
- Excessive bedrest, avoiding activity or movement because of fear of pain.
- Lack of support, overprotective or punitive partner/spouse.
- Impaired sleep or increased alcohol use since onset of pain.

Objectives

- Prevent the progression of acute/subacute to chronic pain through early identification of psychosocial, 'yellow flag', risk factors.
- Provide early multidisciplinary care and develop self-management skills for those who are at risk of transitioning from acute/subacute to chronic pain.
- Minimise biopsychosocial consequence of those who are at risk of chronic pain no matter which intervention is introduced.
- Minimise secondary changes due to the development of chronic pain e.g. physical movement compensation and adaptation, fear avoidance behaviour, depression and anxiety.
- Minimise the burden of healthcare cost of progression to chronic pain and other comorbidities.
- Minimise the overuse of passive modalities of treatment in the early phase of pain management.
- Prevent the overuse of medicines that may result in long term adverse effects (e.g. addiction, dependency, tolerance, endocrine effects).

Service Model

- Similar referral pathway as for the current PainWISE commissioned Turning Pain into Gain Pain Program (TPIG), however the patient referred will be triaged to the acute/subacute pain program.
- Referral form would have a tick box section that identifies the phase of pain experienced by the patient, e.g. 1-3 months post injury/surgery/trauma or 3-6 months post injury/surgery/trauma. Referral form should also include Yellow Flags check list and Orebro Musculoskeletal Pain Screening (10 Questions) – for musculoskeletal pain conditions such as low back pain.
- Education of GPs will also be important and necessary to differentiate the Turning Pain into Gain Persistent Pain Program and the Acute/Subacute pain program. This can be done through GP Newsletter and letters to current network.
- Our triage clinician would act as an early screen for filtering toward the required program. Clear differentiation will be required between the TPIG Persistent Pain program and the Acute/Subacute program.
- Acute/Subacute Self-Management program: A one-day group based education program with up to four 1:1 case management sessions with clinical facilitator (allied health/pharmacist), and 4 allied health sessions as required. Clear hand over to treating allied health clinicians to ensure the same message is reiterated. Final attendance of one-day psychological group based education program prior to discharge. It would be important not to over support. If they need more, they could transition into the TPIG program.
- At discharge a pain management plan will be provided to the patient and communicated to the GP.
- Staggered program start would be necessary to ensure that a new program is commenced once a month. Dedicated staffing required to staff this program to ensure timely access. Waitlist should be no more than 2 weeks.

Target Group/Service Users

- Patients experience pain a) 1-3 months post injury/surgery/trauma or b) 3-6 months post injury/surgery/trauma with 'yellow flag' risk factors indication possible progression to chronic/persistent pain.

Eligibility Criteria:

- The patient has suffered pain 1-3 months post injury/surgery/trauma or for 3-6 months and displaying Yellow Flag indicators suggesting an increased risk of progression to long-term distress, disability and potential drug misuse (the referral will list yellow flag checklist).
- The patient is not displaying any Red Flags (Red Flags are clinical indicators of possible serious underlying conditions requiring further medical intervention).
- The patient is not suitable for surgical or urgent pain specialist interventions.
- The patient is not a palliative care patient.
- The patient requires improved self-management strategies and skills to optimise ongoing care.
- The patient is able to participate in group education.
- Able to give voluntary, informed consent for the ongoing collection of audit data.
- If the patient has had surgery in the past 12 weeks, a functional instructional plan is provided with this referral.

Exclusions:

- Patients who are undergoing worker's compensation.
- This exclusion is recommended for the first year of this pilot study so that the program can be evaluated for its stand alone merits.

Referral Requirements:

- Referral form to be completed with the above *Eligibility Criteria*.
- Referral form will clearly state the name of the program "Early Intervention Acute/ Subacute Pain Program".
- Fax number or Medical Object contact.
- Yellow Flags checklist.
- Confirmation that Red Flags have been excluded.
- Orebro Musculoskeletal Pain Screening (10 Questions).
- Co-referral by Allied Health.
- For Referrals to be valid the patient's GP must sign the referral.
- Mandatory requirements - For patients who have been referred between 6-12 weeks post injury/surgery/trauma, a written instruction of the patient's functional capacity is required from the GP to accompany the referral.

Supporting Collateral Material:

- 3 hour Initial group based "Early Intervention Acute/ Subacute Pain Program" presentation.
- 3 hour Final Psychological group based program presentation.
- Patient Information Sheet specific for the "Early Intervention Acute/ Subacute Pain Program".
- GP Information Sheet specific for the "Early Intervention Acute/ Subacute Pain Program".

Program overview

Total Duration of Acute/Subacute Pain Program = 9 to 12 hours of program involvement over approximately 4 to 6 months

Table 1: Education Program – Duration: 3 hour (10am to 1.00pm or 12.00 to 3.00pm)

40 minutes	What is pain? -Pain and the healing cycle? -Why do I still have pain?
Break 15 mins	
40 minutes	What contributes to my pain?

	<ul style="list-style-type: none"> -Stressors, pain and the brain -Identity and loss -Social engagement and losing your way
Break 15 mins	
40 minutes	<p>Where to from here?</p> <ul style="list-style-type: none"> -Minimising the impact of pain on my life: <ul style="list-style-type: none"> • My mental health • My social engagement -Finding my function: <ul style="list-style-type: none"> • Pacing my functional gains • Thinking outside the box 'what other areas of my body can I engage in with movement?' -Responsible medication usage: <ul style="list-style-type: none"> • Understand the role of medicines in pain management • Understand safety considerations

Table 2: Individual Case Management

Initial assessment	<ul style="list-style-type: none"> • Exclusion of red flags. • Post-surgical patients will need functional instructions from GP of their capacity. • Check current understanding and interpretation of: <p>-Prognosis</p> <p>-Diagnosis</p> <ul style="list-style-type: none"> • Determine and discuss thoughts and actions in relation to the two program goals. • Determine and discuss the barriers of positive outcomes.
Acute/Subacute Self-Management Education Program	<ul style="list-style-type: none"> • Participate in the education program before seeing the clinician for further case management. • 3 hour group based session.
2 nd Case Management Session:	<ul style="list-style-type: none"> • Check current understanding and interpretation of: <p>-Prognosis</p> <p>-Diagnosis</p> <ul style="list-style-type: none"> • Determine and discuss thoughts and actions in relation to two points above. • Determine and discuss the barriers of positive outcomes. • Consider referral to psychologist.
3 rd Case Management Session:	<ul style="list-style-type: none"> • As above. • Navigation to allied health services if needed, 4 extra services above medicare allowance to allied health support if needed.

	<ul style="list-style-type: none"> Consider if referral to longer TPIG program is needed for further support. <p>Or</p> <ul style="list-style-type: none"> Discharge with pain management plan (including a flare up plan and medium term plan).
4 th Case Management Session (optional)	<ul style="list-style-type: none"> If final review is required. Review how allied health or psychologist services are going. Check adherence to plan. Check understanding of pain. Consider if referral to longer TPIG program is needed for further support. <p>OR</p> <ul style="list-style-type: none"> Discharge with pain management plan (including a flare up plan and medium term plan).
<p>Group based Psychological Support Program:</p> <ul style="list-style-type: none"> Summary of key points learned in the Acute/Subacute Self-Management Program. Next steps. Problem solving flare ups. Healthcare navigation. 	<ul style="list-style-type: none"> On exit from the program ALL participants will be invited to a final group based psychological support program. 3 hour group based program. Completion Certificate provided. Discharge with pain management plan (including an Exacerbation 'Flare Up' plan).

<p>Key Outputs: (e.g new service, report etc.)</p> <p>Key Deliverables:</p> <ol style="list-style-type: none"> Increase GP and Allied Health awareness of acute / subacute pain, risk factors and early identification and prevention to chronic pain Enhance collaborations with QLD Health departments and support them in transitioning into primary healthcare for those with acute/subacute pain presenting with Yellow Flag symptoms (e.g. to ED, Post surgery and Rehabilitation) Expand and grow the primary Health Care Network across a community of practice to support community based patients with chronic pain using the consistent messages in promoting self-management and a multimodal model of care. Early referral to other community based social support services Build on existing networks and broaden further collaboration with existing and 	<p>Key Stakeholders: (e.g. GPs, allied health, pharmacist Specialists)</p> <p>GPs, Allied Health, Pharmacists</p> <p>Family, Carers</p> <p>Specialists – public and private</p>
--	--

<p>partners across primary Health care providers.</p> <p>6. A maximum of 40 patients accessing the early intervention acute/subacute pain program between Jan 2020 and June 2020.</p> <p>7. A minimum completion rate of 70% participants (28 participants) within 4-6 months of referral.</p> <p>8. Improved patient experience and quality of care.</p> <p>9. Improved patient, carer and family access to information and education.</p>	
<p>Proposed Start Date: September/ October 2019</p> <p>Timeline of Program Roll out: September and October 2019: Internal operations pathway mapping</p> <p>Nov and Dec 2019: Referral Form completion Patient and GP Information Sheet Completion GP and Allied Health communications</p> <p>January 2020: Patient recruitment commencement</p> <p>February to June 2020: Acute/Subacute Self-Management Program commencement -Group based Acute/Subacute Self-Management Program -Individual case management -Psychological group based support</p>	<p>Estimated Project Duration: Maximum of 6 months</p>
<p>Key risks identified to date:</p> <ul style="list-style-type: none"> • Injury sustained at the pain education program (e.g. falls, setting up, packing up) 	
<p>Project Strategies</p>	
<p>Quality Management (<i>Identify how the quality of the product or service will be assured e.g. peer review, pilots, objective tests, etc.</i>)</p> <ol style="list-style-type: none"> 1. Clinical peer review of content 2. This trial itself is a pilot study of the feasibility of such a program to prevent the progression of acute/subacute pain conditions to chronic/persistent pain. 3. With sufficient funding University Evaluation of the program goals and clinical measures will inform the ongoing quality of the program. 4. Patient feedback utilising both validated questionnaires and pre and post engagement with the program will also help to inform the quality of the program. 	<p>Risk Management - As per current organisation practice.</p>

KPIs:

- A. Early Intervention acute/subacute self-management program referrals – a maximum of 40 patients in 6 months referred
- A. 3 x Acute/Subacute Self-Management Program group presentations
- B. 3 x Psychological Support Program group presentations
- C. Allied Health services- Access to extended allied health interventions (up to 4 extra services per patient).
- D. GP referral numbers into the Early Intervention Acute/Subacute Pain Program
- E. AH referral numbers into the Early Intervention Acute/Subacute Pain Program

COST	
Project Manager 0.1	\$6,000
Project Clinical Co-Facilitator (8 hours/week) @ \$65/hr (for 36 weeks)	\$18,720
Administration Support (8 hours/week) @ \$25/hr (for 36 weeks)	\$7200
Resources : Printing/ mail - @\$40/participant	\$1600
3 x Acute/Subacute Self-Management Program 3 x Psychological Program (venue, catering, resources) approx. \$250/group -Robina Community Centre/ Library	\$1500
Advanced Allied Health Intervention/Therapy (max. 160 sessions) @ \$65/service (Ex GST)	\$10,400
Total	\$ 45,420.00
Evaluation – University <ul style="list-style-type: none"> • ED/Hospital presentation during program • Patient Self-Efficacy Questionnaire • Pain Interference Score pre and post • Orebro pre and post • Change in yellow flag symptoms pre and post 	Approx. \$10,000 (one off for pilot project)
Grand Total	\$ 55,420.00

Signed: _____


—

Name: Joyce McSwan

Position: Clinical Director, PainWISE

Date: 24/8/19

Picture 1: Early Intervention Acute/Subacute Self-Management Pain Program Pathway



References for Project Plan

- Glattacker, M., Heyduck, K., & Jakob, T. (2018). Yellow flags as predictors of rehabilitation outcome in chronic low back pain. *Rehabilitation Psychology, 63*(3), 408-417. doi: 10.1037/rep0000200
- Hunter New England Health. (2019). Recovery from pain. Retrieved 24 August 2019, from <http://www.hnehealth.nsw.gov.au/Pain/Pages/Pain.aspx>
- Main, C. (2013). The importance of psychosocial influences on chronic pain. *Pain Management, 3*(6), 455-466. doi: 10.2217/pmt.13.49
- Main, C., Kendall, N., & Hasenbring, M. (2012). Screening of Psychosocial Risk Factors (Yellow Flags) for Chronic Back Pain and Disability. *From Acute To Chronic Back Pain*, 203-229. doi: 10.1093/med/9780199558902.003.0060
- Nicholas, M., Linton, S., Watson, P., & Main, C. (2011). Early Identification and Management of Psychological Risk Factors ("Yellow Flags") in Patients With Low Back Pain: A Reappraisal. *Physical Therapy, 91*(5), 737-753. doi: 10.2522/ptj.20100224
- Pain Management Network. (2019). Chronic Pain. Retrieved 24 August 2019, from <https://www.aci.health.nsw.gov.au/chronic-pain>
- Sullivan, M., Feuerstein, M., Gatchel, R., Linton, S., & Pransky, G. (2005). Integrating Psychosocial and Behavioral Interventions to Achieve Optimal Rehabilitation Outcomes. *Journal Of Occupational Rehabilitation, 15*(4), 475-489. doi: 10.1007/s10926-005-8029-9

Appendix 10: Key content of the training packages in the IMPaCT Back trial

Sowden et al (66)

Key content covered in the training packages.

	Key content covered
GP best practice updates	<p>The subgrouping and targeted treatment system and study design, protocols and relevant documentation</p> <p>Use and interpretation of the 6-item subgrouping tool to guide treatment and referral to physiotherapy</p> <p>Screening for red flags and diagnosis</p> <p>Reassurance about good overall prognosis, the benign nature of the LBP and addressing concerns</p> <p>Role of further investigations</p> <p>Simple messages and advice about pain medication</p> <p>Appropriate use of pain relieving modalities</p> <p>Advice about appropriate physical activity levels, return to normal activity, including work and avoiding bed rest</p> <p>Sickness certification</p>
Low- and medium-risk training	<p>The subgrouping and targeted treatment system and study design, protocols and relevant documentation</p> <p>Use and interpretation of the subgrouping tool to guide treatment</p> <p>The role of diagnostic investigations, medication, epidural injections and surgery in back pain and radiculopathy</p> <p>Appropriate reassurance and explanation re: low back pain symptoms</p> <p>Appropriate advice about analgesia.</p> <p>Advice about the maintenance of, or return to, usual activities (including work)</p> <p>Onwards referral of patients who present a clinical or management concern (e.g. those with signs of potential serious pathology or red flags or significant radicular symptoms)</p> <p>Current guidelines for managing LBP in primary care, including current best physiotherapy practice for the management of disability, back pain and referred leg pain, including the role of exercise and manual therapy as well as strategies for equipping patients with the skills to manage future recurrences.</p> <p>Goal setting, pacing and graded exercise will be covered briefly</p> <p>The configuration and availability of local services such as interface clinics and secondary care spinal services and how to refer study patients to these services</p> <p>The role of Job Centre Plus in return to work facilitation.</p>
High-risk training	<p>Specific biopsychosocial factors that contribute to the development and maintenance of chronic pain and disability</p> <p>The importance of key processes and how to utilise knowledge of them in treatment</p> <p>Identifying key psychosocial prognostic indicators using stem and leaf questions</p> <p>Exploring the impact of an individuals' pain on activity, work, sleep, relationships and mood</p> <p>Basic and advanced communications skills training including rapport building, listening, demonstrating empathy and motivational interviewing skills</p> <p>Facilitating discussions with patients about the relation between physical and psychosocial factors</p> <p>Applying the biopsychosocial and cognitive behavioural models to the management of pain and pain related disability and distress</p> <p>Making sense of the assessment information, clinical reasoning, identifying appropriate targets for treatment and treatment planning</p> <p>Explaining pain and providing reassurance</p> <p>Problem solving difficulties</p> <p>Managing patients' expectations</p> <p>Promoting an active rehabilitation self-management approach</p> <p>Challenge patients' unhelpful or inaccurate beliefs and expectations, for example through the provision of individualized information, reassurance and advice</p> <p>Pacing and graded activity in order to sustain or increase meaningful physical function</p> <p>Improving sleep, mood, social and work functioning</p> <p>Dealing with distressed or complex patients and when to refer onwards or seek additional input</p> <p>Goal setting</p> <p>Monitoring and reinforcing progress and modifying treatment</p> <p>Supporting patients in active self-management of future set-backs or recurrences</p> <p>How to appropriately refer or signpost patients to other services such as chronic pain services</p>

The 9-item STarT Back screening tool.

Patient name: _____ Date: _____

Thinking about the last 2 weeks tick your response to the following questions:

	Disagree 0	Agree 1
1 My back pain has spread down my leg(s) at some time in the last 2 weeks	<input type="checkbox"/>	<input type="checkbox"/>
2 I have had pain in the shoulder or neck at some time in the last 2 weeks	<input type="checkbox"/>	<input type="checkbox"/>
3 I have only walked short distances because of my back pain	<input type="checkbox"/>	<input type="checkbox"/>
4 In the last 2 weeks, I have dressed more slowly than usual because of back pain	<input type="checkbox"/>	<input type="checkbox"/>
5 It's not really safe for a person with a condition like mine to be physically active	<input type="checkbox"/>	<input type="checkbox"/>
6 Worrying thoughts have been going through my mind a lot of the time	<input type="checkbox"/>	<input type="checkbox"/>
7 I feel that my back pain is terrible and it's never going to get any better	<input type="checkbox"/>	<input type="checkbox"/>
8 In general I have not enjoyed all the things I used to enjoy	<input type="checkbox"/>	<input type="checkbox"/>
9 Overall, how bothersome has your back pain been in the last 2 weeks?		

Not at all <input type="checkbox"/> 0	Slightly <input type="checkbox"/> 0	Moderately <input type="checkbox"/> 0	Very much <input type="checkbox"/> 1	Extremely <input type="checkbox"/> 1
---	---	---	--	--

Total score (all 9): _____ **Sub Score (Q5-9):** _____

© Keele University 01/08/07

Appendix 11: Content of a GP trainee education initiative about opioids implemented in Australia

Holliday (110)

Box 1 Presentation content

The history of opium and analgesia practice

The escalation in the West of opioid prescribing and associated harms, including overdose and addiction

Chronic noncancer pain (CNCP) neurophysiology including neuro-plasticity, central sensitization, and opioid-induced hyperalgesia

Guideline-concordant and patient-centred management of CNCP

Biopsychosocial assessment in CNCP including past and present psychiatric and substance use problems, in preference to tool-based risk stratification (38)

Use of the Pain Intensity, Enjoyment of Life, General Activity measurement scale (40)

The importance of multidisciplinary and multimodal CNCP management with appropriate referral to physiotherapy, psychology, pain specialists, or addiction treatment services

The nonpharmaceutical self-management management of CNCP

The nonopioid pharmaceutical management of CNCP

The lack of evidence supporting opioids in CNCP in terms of efficacy and safety

The practice, principals, and limitations of universal precautions if or when opioids are used in CNCP

The importance of assessing and responding to the emergence of aberrant behavior

Deprescribing opioids