

What large-scale lifestyle interventions work to prevent type 2 diabetes?

Background

- More than a million Australians have diabetes (all types) or are at high risk of diabetes. Type 2 diabetes makes up around 85% of these diabetes cases.
- Progression to diabetes can be prevented or delayed. Modifiable risk/protective factors for diabetes include body mass index, physical activity, smoking status and early life factors such as birthweight.
- Many lifestyle interventions for diabetes prevention help to prevent other chronic diseases such as cardiovascular disease.
- Delaying the onset and severity of diabetes is important because it can prevent diabetes-related complications.
- There have been at least 13 large-scale lifestyle intervention trials, and some additional associated follow-up and outcome studies. These interventions have taken place in Australia^{1,2}, China³, Finland⁴⁻⁷, Germany⁸, India⁶, Japan^{6,9}, the United Kingdom⁸ and the United States¹⁰.

Review purpose

- To summarise existing evidence regarding type 2 diabetes prevention interventions, focusing on lifestyle interventions, and to showcase the extent to which different lifestyle interventions have been effective.
- To assess the effect size of different lifestyle interventions, and to consider the potential for scalability.
- To identify research gaps and areas of most potential to inform future research investment.

Key findings

- Twelve papers met the selection criteria. All the interventions demonstrated an effect on diabetes risk reduction.
- Some seemingly similar interventions had different effect sizes in different studies. Effect size differed according to the length of the intervention (tended to increase with length), length of follow-up (tended to decrease with length) and participation levels (tended to increase with high participation).
- While not lifestyle interventions, bariatric surgery and medications were included as comparators in many studies assessing lifestyle interventions. For this reason, bariatric surgery and medications were also included in this review.
- Bariatric surgery was the only intervention with a large effect size, but the studies only assessed participants who were obese and at high risk. Bariatric surgery is likely to be inapplicable and unfeasible at the general population level.
- Interventions with a medium effect size were: diet and/or exercise; diet and exercise and an additional component such as education; and exercise and/or education.
- Not all existing lifestyle intervention studies were included in this review as they did not meet the inclusion criteria. Many population-level interventions measured weight loss or other clinical indicators as a proxy for impact on diabetes risk¹¹⁻¹⁷. Many study participants had successful and sustained weight loss, suggesting reduced diabetes risk. However, this information was inadequate to confidently calculate effect size.

Summary map of evidence

	Very small* – small effect odds ratio (OR) 1.5-2, relative risk (RR) 2-3	Medium effect OR 2-3, RR 3-4	Large effect OR > 3, RR > 4
Diet	n=1172 MA ¹⁸		
Diet +/- exercise	n=400 Da Qing ^{3,19} ★ n=147 SLIM ⁶	n=2993 MA ²⁰ n=4581 MA ¹⁸ n=458 Japan ⁶ ★	
Diet + education**	n=641 Zensharen ⁹ ★		
Diet + exercise + other	n=655 MA ²¹ n=531 IDPP ⁶ ★ n=1079 DPP ¹⁰ ★ n=910 DPPOS ²² ★	n=194 Sweden ⁶	
Exercise +/- education**	n=52,475 MA ¹⁸ n=522 DPS ^{6,23} ★	n=2798 FIN-D2D ^{4,24} ★	
Anti-diabetic drugs (Metformin)	n=18,155 MA ²⁰ n=7149 MA ¹⁸ n=1073 DPP ¹⁰ ★ n=942 DPPOS ²² ★		
Bariatric surgery			n=3650 MA ²⁰ n=9400 MA ¹⁸

LEGEND

★ = Single, large-scale trial (n ≥ 400)

Systematic review

RCT

KEY TO SUMMARY MAP

- Da Qing, Da Qing Study, China
- DPP, Diabetes Prevention Program, the United States
- DPPOS, Diabetes Prevention Program Outcomes Study, the United States
- DPS, Diabetes Prevention Study, Finland
- FIN-D2D, National Type 2 Diabetes Prevention Program, Finland
- IDPP, Indian Diabetes Prevention Program, India
- Japan, randomised controlled trial in Japan
- MA, meta-analysis, value from more than one study
- SLIM, Study on Lifestyle intervention and Impaired glucose tolerance Maastricht, Netherlands
- Sweden, randomised controlled trial in Sweden
- Zensharen, Zensharen Study, Japan.

* Very small includes interventions with OR < 1.5 and RR < 2 if study authors stated that an intervention was significant.

** Education includes lectures, educational material and individual or group counselling.

Research gaps

- More research is needed to determine what 'dose' or intensity of lifestyle intervention is needed, what components of programs are most effective, and how programs should be modified to target specific population groups.
- There is little published evidence about the successful scaling up of interventions. Better research on the mainstreaming of clinical trial size interventions to population intervention programs is needed.
- Some interventions with the same components (e.g. diet and/or exercise) are more effective than others. Determining what characteristics make an intervention effective could lead to greater effect sizes of these interventions, and greater potential for successful scaling up.
- While there is good evidence that type 2 diabetes can be prevented or delayed, further research is needed to assess the effect size of lifestyle interventions at the general population level. Many existing lifestyle intervention studies did not assess diabetes incidence, compare the intervention group against a control group, or have not published results.

Summary of review method

- An electronic search was conducted of relevant articles published to March 2015, identified from the electronic databases including PubMed, Scopus and Google Scholar. Seminal articles were also identified through consultation with experts in the field.
- Studies were included if they measured diabetes incidence as an outcome and had an intervention duration of at least six months. Some large-scale type 2 diabetes interventions measured weight change and other clinical measurements but not diabetes incidence. Even though some of these other outcomes are known to be protective against diabetes, these studies were excluded because their effect size for reducing type 2 diabetes could not be confidently determined with confidence. To capture a breadth of population-wide interventions, all controlled study designs were included. As the scope of the review was primary prevention of type 2 diabetes, study participants were required to not have diabetes at baseline. Where possible, trials were only included once, and studies were excluded if they only offered duplicate data.
- Trials were grouped according to intervention domain (e.g. diet, diet and/or exercise, exercise and/or education), with some trials having multiple domains due to multiple cohorts. Data were extracted on study design, study population, intervention characteristics and primary outcomes (prevention of diabetes).
- A guide²⁵ was used to aid the classification of an intervention's effect size. If an effect was classified as no effect using the guide, but the authors of the study had reported an effect, a small effect size was given to the intervention. Where multiple effect sizes were measured during a study, the peak effect size was used to demonstrate what effect was possible for that intervention.
- **Limitations:** A rapid review approach was used. Completeness of such reviews are determined by time constraints. Potential bias associated with lack of inclusion of relevant studies was minimised by reliance on recent comprehensive systematic reviews, consultation with experts in the field, and an iterative snowball approach to identify any additional relevant papers from reference lists. In the absence of detailed data from each study, effect sizes were calculated on the basis of what was reported in studies. While every effort was made to ensure consistency when analysing studies' effect sizes, the heterogeneity of studies meant that comparisons between interventions can only be taken as a guide. For example, some of the studies were conducted only in males or in people that already had impaired glucose tolerance at baseline. However, this is not a significant limitation for the purpose of showcasing different types of interventions and their potential to prevent diabetes.

The Australian Prevention Partnership Centre

The Prevention Centre is finding out how we can build an effective, efficient and equitable system for the prevention of lifestyle-related chronic disease.

This report has been prepared by Emily Morrice, Dr Jo-An Atkinson and Professor Andrew Wilson.

© Sax Institute 2015



Contact us:

Tel: (02) 9188 9520

Email: preventioncentre@saxinstitute.org.au

Website: preventioncentre.org.au

References

- Colagiuri S, Vita P, Cardona-Morrell M et al. The Sydney Diabetes Prevention Program: a community-based translational study. *BMC Public Health* 2010;10:328.
- Dunbar JA, Jayawardena A, Johnson G et al. Scaling up diabetes prevention in Victoria, Australia: policy development, implementation and evaluation. *Diabetes Care* 2014;37(4):934–42.
- Pan XR, Li GW, Hu YH et al. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 1997;20(4):537–44.
- Saaristo T, Moilanen L, Korpi-Hyovalti E et al. Lifestyle intervention for prevention of type 2 diabetes in primary health care: one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D). *Diabetes Care* 2010;33(10):2146–51.
- Absetz P, Valve R, Oldenburg B et al. Type 2 diabetes prevention in the “real world”: One-year results of the GOAL Implementation Trial. *Diabetes Care* 2007;30(10):2465–70.
- Yoon U, Kwok LL, Magkidis A. Efficacy of lifestyle interventions in reducing diabetes incidence in patients with impaired glucose tolerance: a systematic review of randomized controlled trials. *Metabolism* 2013;62(2):303–14.
- Viitasalo K, Hemiö K, Puttonen S, et al. Prevention of diabetes and cardiovascular diseases in occupational health care: Feasibility and effectiveness. *Prim Care Diabetes* 2014;9(2):96–104.
- Schwarz PE. Public health implications: translation into diabetes prevention initiatives – four-level public health concept. *Med Clin North Am* 2011;95(2):397–407, ix.
- Saito T, Watanabe M, Nishida J et al. Lifestyle modification and prevention of type 2 diabetes in overweight Japanese with impaired fasting glucose levels: a randomized controlled trial. *Arch Intern Med* 2011;171(15):1352–60.
- Knowler WC, Barrett-Connor E, Fowler SE et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346(6):393–403.
- Schwarz P, Reddy P. *Prevention of Diabetes*. UK: John Wiley & Sons Ltd. 2013.
- Salopuro TM, Saaristo T, Oksa H, et al. Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity. *BMC Public Health* 2011;11:350.
- Dunbar J, Colagiuri S, Reddy P, et al. Scaling up type 2 diabetes prevention programs: National and State interventions in Australia. *Diabetes Prevention in Practice*. Dresden: Tumaini Institute from Prevention Management; 2010:45–56.
- Makrilakis K, Liatis S, Grammatikou S, Perrea D, Katsilambros N. Implementation and effectiveness of the first community lifestyle intervention programme to prevent Type 2 diabetes in Greece. The DE-PLAN study. *Diabet Med* 2010;27(4):459–65.
- Dunkley AJ, Bodicoat DH, Greaves CJ, et al. Diabetes prevention in the real world: effectiveness of pragmatic lifestyle interventions for the prevention of type 2 diabetes and of the impact of adherence to guideline recommendations: a systematic review and meta-analysis. *Diabetes Care* 2014;37(4):922–33.
- Angermayr L, Melchart D, Linde K. Multifactorial lifestyle interventions in the primary and secondary prevention of cardiovascular disease and type 2 diabetes mellitus – a systematic review of randomized controlled trials. *Ann Beh Med* 2010;40(1):49–64.
- Ali MK, Echouffo-Tcheugui J, Williamson DF. How effective were lifestyle interventions in real-world settings that were modeled on the Diabetes Prevention Program? *Health Aff* 2012;31(1):67–75.
- Merlotti C, Morabito A, Pontiroli AE. Prevention of type 2 diabetes; a systematic review and meta-analysis of different intervention strategies. *Diabetes Obes Metab* 2014;16(8):719–27.
- Li G, Zhang P, Wang J, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 2008;371(9626):1783–9.
- Merlotti C, Morabito A, Ceriani V, Pontiroli AE. Prevention of type 2 diabetes in obese at-risk subjects: a systematic review and meta-analysis. *Acta Diabetol* 2014; Oct;51(5):853–63.
- Schellenberg ES, Dryden DM, Vandermeer B, Ha C, Korownyk C. Lifestyle interventions for patients with and at risk for type 2 diabetes: a systematic review and meta-analysis. *Ann Intern Med* 2013;159(8):543–51.
- Knowler WC, Fowler SE, Hamman RF, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374(9702):1677–86.
- Lindstrom J, Peltonen M, Eriksson JG, et al. Improved lifestyle and decreased diabetes risk over 13 years: long-term follow-up of the randomised Finnish Diabetes Prevention Study (DPS). *Diabetologia* 2013;56(2):284–93.
- Saaristo T, Peltonen M, Keinänen-Kiukkaanniemi S, et al. National type 2 diabetes prevention programme in Finland: FIN-D2D. *Int J Circumpolar Health* 2007;66(2):101–12.
- Sullivan GM, Feinn R. Using effect size – or why the P value is not enough. *J Grad Med Educ* 2012;4(3):279–82.

Our funding partners



Co-hosted by

